

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07K 14/00</b>	<b>A2</b>	(11) International Publication Number: <b>WO 99/33868</b> (43) International Publication Date: 8 July 1999 (08.07.99)
<p>(21) International Application Number: PCT/EP98/08563</p> <p>(22) International Filing Date: 18 December 1998 (18.12.98)</p> <p>(30) Priority Data: 9727262.9 24 December 1997 (24.12.97) GB</p> <p>(71) Applicant (for all designated States except US): SMITHKLINE BEECHAM BIOLOGICALS S.A. [BE/BE]; Rue de l'Institut 89, B-1330 Rixensart (BE).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): DALEMANS, Wilfried, L., J. [BE/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE). GERARD, Catherine, Marie, Ghislaine [BE/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE).</p> <p>(74) Agent: DALTON, Marcus, Jonathan, William; SmithKline Beecham, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p>
<p>(54) Title: VACCINE</p> <p>(57) Abstract</p> <p>The present invention provides Human Papilloma Virus (HPV) fusion proteins, linked to an immunological fusion partner that provides T helper epitopes to the HPV antigen. Vaccine formulations are provided that are useful in the treatment or Prophylaxis of HPV induced tumours.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## VACCINE

The present invention relates to vaccine compositions, comprising an E6 or/ and E7 or E6, E7 fusion protein from an HPV strain optionally linked with an immunological fusion partner and formulated with a CpG containing oligonucleotide  
5 into vaccines that find utility in the treatment or prophylaxis of human papilloma virus induced tumours or lesions. In particular, the present invention relates to vaccines comprising fusions proteins, comprising a protein or part of a protein that provides T helper epitopes (such as protein D from Haemophilus influenzae B) and an antigen  
10 from a human-papilloma virus (eg comprising an E6 or E7 protein from HPV 16 or 18 strain associated with cancer) that find utility in the treatment or prophylaxis of human papilloma induced tumours, wherein the vaccine is formulated with a CpG containing oligonucleotide as an adjuvant.

Papillomaviruses are small naked DNA tumour viruses (7.9 kilobases, double  
15 strand), which are highly species-specific. Over 70 individual human papillomavirus (HPV) genotypes have been described. Papillomaviruses are classified on the basis of species of origin (human, bovine etc.) and of the degree of genetic relatedness with other papillomaviruses from the same species. HPVs are generally specific for the skin or mucosal surfaces and have been broadly classified into "low" and "high" risk  
20 viruses.

Low risk HPVs usually cause benign *lesions* (warts or papillomas) that persist for several months or years. High risk HPVs are associated with pre-neoplastic lesions and cancer. The strongest positive association between an HPV virus and human cancer is that which exist between HPV 16 and 18 and cervical carcinoma.  
25 More than ten other HPV types have also been found in cervical carcinomas including HPV 31 and HPV 33 although at less frequency.

Genital HPV infection in young sexually active women is common and most individuals either clear the infection, or if lesions develop, these regress. Only a subset of infected individuals has lesions which progress to high grade intraepithelial  
30 neoplasia and only a fraction of these progress further to invasive carcinoma.

The molecular events leading to HPV infection have not been clearly established. The lack of an adequate *in vitro* system to propagate human

papillomaviruses has hampered the progress to a best information about the viral cycle.

Today, the different types of HPVs have been isolated and characterised with the help of cloning systems in bacteria and more recently by PCR amplification. The molecular organisation of the HPV genomes has been defined on a comparative basis with that of the well characterised bovine papillomavirus type 1 (BPV1).

Although minor variations do occur, all HPVs genomes described have at least seven early genes, E1 to E7 and two late genes L1 and L2. In addition, an upstream regulatory region harbors the regulatory sequences which appears to control most transcriptional events of the HPV genome.

E1 and E2 genes are involved in viral replication and transcriptional control, respectively and tend to be disrupted by viral integration. E6 and E7 are involved in viral transformation. E5 has also been implicated in this process.

In the HPVs involved in cervical carcinoma such as HPV 16 and 18, the oncogenic process starts after integration of viral DNA. The integration results in the inactivation of genes coding for the capsid proteins L1 and L2 and loss of E2 repressor function leads to deregulation of the E6/E7 open reading frame installing continuously overexpression of the two early proteins E6 and E7 that will lead to gradually loss of the normal cellular differentiation and the development of the carcinoma. E6 and E7 overcome normal cell cycle by inactivating major tumor suppressor proteins, p53 and pRB, the retinoblastoma gene product, respectively.

Carcinoma of the cervix is common in women and develops through a pre-cancerous intermediate stage to the invasive carcinoma which frequently leads to death. The intermediate stages of the disease is known as cervical intraepithelial neoplasia and is graded I to III in terms of increasing severity (*CIN I-III*).

Clinically, HPV infection of the female anogenital tract manifests as cervical flat condylomas, the hallmark of which is the koilocytosis affecting predominantly the superficial and intermediate cells of the cervical squamous epithelium.

Koilocytes which are the consequence of a cytopathic effect of the virus, appear as multinucleated cells with a perinuclear clear haloe. The epithelium is thickened with abnormal keratinisation responsible for the warty appearance of the lesion.

Such flat condylomas when positive for the HPV 16 or 18 serotypes, are high-risk factors for the evolution toward cervical intraepithelial neoplasia (CIN) and carcinoma in situ (CIS) which are themselves regarded as precursor lesions of invasive cervix carcinoma.

5           The natural history of oncogenic HPV infection presents three consecutive phases, namely:

- (1) a latent infection phase,
- (2) a phase of intranuclear viral replication with product of complete virions, which corresponds to the occurrence of koilocytes. At this stage, the HPV is producing its  
10 full range of proteins including E2, E5, E6, E7, L1 and L2.
- (3) a phase of viral integration into the cellular genome, which triggers the onset of malignant transformation, and corresponds to CIN II and CIN III/CIS with progressive disappearance of koilocytes. At this stage, the expression of E2 is down-regulated, the expression of E6 and E7 is enhanced. Between CIN II/III and CIN III /  
15 Cervix carcinoma the viral DNA changes from being episomal in the basal cells to integration of E6 and E7 genes only (tumoral cells). 85% of all cervix carcinomas are squamous cell carcinomas most predominantly related to the HPV16 serotype. 10% and 5% are adenocarcinomas and adenosquamous cell carcinomas respectively, and both types are predominantly related to HPV 18 serotype. Nevertheless other  
20 oncogenic HPV's exist.

International Patent Application No. WO 96/19496 discloses variants of human papilloma virus E6 and E7 proteins, particularly fusion proteins of E6/E7 with a deletion in both the E6 and E7 proteins. These deletion fusion proteins are said to be immunogenic.

25           Immunomodulatory oligonucleotides contain unmethylated CpG dinucleotides ("CpG") and are known (WO 96/02555, EP 468520). CpG is an abbreviation for cytosine-guanosine dinucleotide motifs present in DNA. Historically, it was observed that the DNA fraction of BCG could exert an anti-tumour effect. In further studies, synthetic oligonucleotides derived from BCG gene sequences were shown to be  
30 capable of inducing immunostimulatory effects (both in vitro and in vivo). The authors of these studies concluded that certain palindromic sequences, including a central CG motif, carried this activity. The central role of the CG motif in

immunostimulation was later elucidated in a publication by Krieg, Nature 374, p546 1995. Detailed analysis has shown that the CG motif has to be in a certain sequence context, and that such sequences are common in bacterial DNA but are rare in vertebrate DNA.

5 It is currently believed that this evolutionary difference allows the vertebrate immune system to detect the presence of bacterial DNA (as occurring during an infection) leading consequently to the stimulation of the immune system. The immunostimulatory sequence as defined by Krieg is:

Purine Purine CG pyrimidine pyrimidine and where the CG motif is not  
10 methylated. In certain combinations of the six nucleotides a palindromic sequence is present. Several of these motifs, either as repeats of one motif or a combination of different motifs, can be present in the same oligonucleotide. The presence of one or more of these immunostimulatory sequence containing oligonucleotides can activate various immune subsets, including natural killer cells (which produce interferon  $\gamma$  and  
15 have cytolytic activity) and macrophages (Wooldridge et al Vol 89 (no. 8), 1977). Although other unmethylated CpG containing sequences not having this consensus sequence have now been shown to be immunomodulatory.

The present invention provides compositions comprising either an E6 or/and E7 or an E6/E7 fusion protein optionally linked to an immunological fusion partner  
20 having T cell epitopes, and adjuvanted with an immunomodulatory CpG containing oligonucleotide.

In a preferred form of the invention, the immunological fusion partner is derived from protein D of Haemophilus influenza B. Preferably the protein D derivative comprises approximately the first 1/3 of the protein, in particular  
25 approximately the first N-terminal 100-110 amino acids. The protein D may be lipidated (Lipo Protein D). Other immunological fusion partners include the non-structural protein from influenzae virus, NS1 (hemagglutinin). Typically the N terminal 81 amino acids are utilised, although different fragments may be used provided they include T-helper epitopes.

30 In another embodiment the immunological fusion partner is the protein known as LYTA. Preferably the C terminal portion of the molecule is used. Lyta is derived from Streptococcus pneumoniae which synthesize an N-acetyl-L-alanine amidase,

amidase LYTA, (coded by the *lytA* gen {Gene, 43 (1986) page 265-272} an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-terminal domain of the LYTA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of E.coli C-LYTA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LYTA fragment at its amino terminus has been described {Biotechnology: 10, (1992) page 795-798}. As used herein a preferred embodiment utilises the repeat portion of the Lyta molecule found in the C terminal end starting at residue 178. A particularly preferred form incorporates residues 188 - 305.

Accordingly, the present invention in preferred embodiment provides compositions comprising an immunomodulatory CpG oligonucleotide and a fusion proteins comprising Protein D - E6 from HPV 16, Protein D - E7 from HPV 16 Protein D - E7 from HPV 18, Protein D - E6 from HPV 18, and Protein D E6 E7 from both HPV 16 and 18. The protein D part preferably comprises the first 1/3 of protein D. It will be appreciated that other E6 and E7 proteins may be utilised from other HPV subtypes.

The proteins utilised in the present invention preferably are expressed in E. coli. In a preferred embodiment the proteins are expressed with a Histidine tail comprising between 5 to 9 and preferably six Histidine residues. These are advantageous in aiding purification.

The protein E7 may in a preferred embodiment carry one or several mutations in the binding site for the *rb* (retinoblastoma gene product) and hence eliminate any potential transforming capacity. Preferred mutations for HPV 16 E7 involve replacing Cys<sub>24</sub> with Glycine, or Glutamic acid<sub>26</sub> with Glutamine. In a preferred embodiment the E7 protein contains both these mutations.

Preferred mutations for the HPV 18 E<sub>7</sub> involve replacing Cys<sub>27</sub> with Glycine and/or Glutamic acid<sub>29</sub> with Glutamine. Again preferably both mutations are present.

Single or double mutations may also be introduced p53 region of E<sub>6</sub> to eliminate any potential transforming ability.

In a further embodiment of the invention there is provided an E6 E7 fusion protein from HPV linked to an immunological fusion partner and a CpG immunomodulatory oligonucleotide.

The vaccine of the present invention preferentially induce a TH1 immune  
5 response.

Two main types of Helper T cells have been characterized TH1 and TH2, which differ in the type of cytokines they secrete. These cytokines can be considered as the driving force behind the development of 2 different types of immune response : TH1-type of immune response is associated with cell mediated effector mechanisms  
10 such as production of the INF- $\gamma$  and IL-2 cytokines by T-lymphocytes. INF- $\gamma$  which in turn can activate other cells and induce them to secrete other important cytokines and mediators (INF- $\gamma$  - activated NK cells produce IL12, IL2-activated NK cells are transformed into lymphokine activated killer cell (LAK), INF- $\gamma$ -activated  
macrophages secrete inflammatory mediators like TNF $\alpha$ , IL1, IL6 and release nitric  
15 oxyde, IL2 can provide help for the differentiation of antigen specific, haplotype restricted cytotoxic T lymphocytes (CTL). At the antibody level, in mice, Th1-type of immune response is also associated with the generation of antibodies of the IgG2 isotype (IgG2a in Balb/c mice and IgG2b in C57BL/6 mice).

The Th2-type of immune response is associated with a humoral immune  
20 response to the antigen, with the production of cytokines like IL4, IL5, IL6, IL10 and by the generation of a broad range of immunoglobulin isotypes including in mice IgG1, IgA, and IgM.

In man the distinction of Th1 and Th2-type immune responses is not absolute. An individual will support an immune response which is predominantly Th1 or  
25 predominantly Th2. However, it is often convenient to consider the families of cytokines in terms of that described in murine CD4 +ve T cell clones by Mosmann and Coffman (*Mosmann, T.R. and Coffman, R.L. (1989) TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. Annual Review of Immunology, 7, p145-173*).

30 In the human TH1 type of response is also associated with the presence of cytokine (IFN $\gamma$  and IL2) eventually with the presence of CTL and IgG2 isotypes in mice correspond to IgG1 type antibodies



This type 1 phenotype is of particular importance in protecting against viral and intracellular bacterial infections as well as in the treatment of cancer.

To manufacture the proteins used in the invention by recombinant techniques, an expression strategy can be used which involves fusion of E7, E6 or E6/E7 fusion to the 1/3-N-terminal portion of protein D from Haemophilus influenzae B, an immunological fusion partner providing T cell helper epitopes. An affinity polyhistidine tail is engineered at the carboxy terminus of the fusion protein allowing for simplified purification. Such recombinant antigen is overexpressed in E. coli as insoluble protein.

The proteins of the invention may be coexpressed with thioredoxin in trans (TIT). Coexpression of thioredoxin in trans versus in cis is preferred to keep antigen free of thioredoxin without the need for protease. Thioredoxin coexpression eases the solubilisation of the proteins of the invention. Thioredoxin coexpression has also a significant impact on protein purification yield, on purified-protein solubility and quality.

The replicable expression vectors may be prepared in accordance with the invention, by cleaving a vector compatible with the host cell to provide a linear DNA segment having an intact replicon, and combining said linear segment with one or more DNA molecules which, together with said linear segment encode the desired product, such as the DNA polymer encoding the protein of the invention, or derivative thereof, under ligating conditions.

Thus, the DNA polymer may be preformed or formed during the construction of the vector, as desired.

The choice of vector will be determined in part by the host cell, which may be prokaryotic or eukaryotic but preferably is E. coli. Suitable vectors include plasmids, bacteriophages, cosmids and recombinant viruses.

The preparation of the replicable expression vector may be carried out conventionally with appropriate enzymes for restriction, polymerisation and ligation of the DNA, by procedures described in, for example, Maniatis *et al.* cited above.

The recombinant host cell is prepared, in accordance with the invention, by transforming a host cell with a replicable expression vector of the invention under transforming conditions. Suitable transforming conditions are conventional and are

described in, for example, Maniatis *et al.* cited above, or "DNA Cloning" Vol. II, D.M. Glover ed., IRL Press Ltd, 1985.

The choice of transforming conditions is determined by the host cell. Thus, a bacterial host such as *E. coli* may be treated with a solution of  $\text{CaCl}_2$  (Cohen *et al.*, Proc. Nat. Acad. Sci., 1973, 69, 2110) or with a solution comprising a mixture of RbCl,  $\text{MnCl}_2$ , potassium acetate and glycerol, and then with 3-[N-morpholino]-propane-sulphonic acid, RbCl and glycerol. Mammalian cells in culture may be transformed by calcium co-precipitation of the vector DNA onto the cells. The invention also extends to a host cell transformed with a replicable expression vector of the invention.

Culturing the transformed host cell under conditions permitting expression of the DNA polymer is carried out conventionally, as described in, for example, Maniatis *et al.* and "DNA Cloning" cited above. Thus, preferably the cell is supplied with nutrient and cultured at a temperature below  $50^\circ\text{C}$ .

The product is recovered by conventional methods according to the host cell. Thus, where the host cell is bacterial, such as *E. coli* it may be lysed physically, chemically or enzymatically and the protein product isolated from the resulting lysate. Where the host cell is mammalian, the product may generally be isolated from the nutrient medium or from cell free extracts. Conventional protein isolation techniques include selective precipitation, adsorption chromatography, and affinity chromatography including a monoclonal antibody affinity column.

When the proteins of the present invention are expressed with a histidine tail (His tag). The proteins can easily be purified by affinity chromatography using an ion metal affinity chromatography column (IMAC) column.

A second chromatographic step, such as Q-sepharose may be utilised either before or after the IMAC column to yield highly purified protein. If the immunological fusion partner is C-LYTA, then it is possible to exploit the affinity of CLYTA for choline and/or DEAE to purify this product. Products containing both C-LYTA and his tags can be easily and efficiently purified in a two step process involving differential affinity chromatography. One step involves the affinity of the His tag to IMAC columns, the other involves the affinity of the C-terminal domain of LYTA for choline or DEAE.

A preferred vaccine composition comprises at least Protein D - E6 from HPV 16 or derivative thereof together with Protein D - E7 from HPV 16. Alternatively the E6 and E7 may be presented in a single molecule, preferably a Protein D E6/E7 fusion. Such vaccine may optionally contain either or both E6 and E7 proteins from HPV 18, preferably in the form of a Protein D - E6 or Protein D - E7 fusion protein or Protein D E6/E7 fusion protein. The vaccines of the present invention may contain other HPV antigens from HPV 16 or 18. In particular, the vaccine may contain L1 or L2 antigen monomers. Alternatively such L1 or L2 antigens may be presented together as a virus like particle or the L1 alone protein may be presented as virus like particle or capsomer structure. Such antigens, virus like particles and capsomer are per se known. See for example WO94/00152, WO94/20137, WO94/05792, and WO93/02184. Additional early proteins may be included such as E2 or preferably E5 for example. The vaccine of the present invention may additionally comprise antigens from other HPV strains, preferably from strains HPV 6, 11, 31 or 33.

Vaccine preparation is generally described in Vaccine Design - The subunit and adjuvant approach (Ed. Powell and Newman) Pharmaceutical Biotechnology Vol. 6 Plenum Press 1995. Encapsulation within liposomes is described by Fullerton, US Patent 4,235,877.

The preferred oligonucleotides preferably contain two or more CpG motifs separated by six or more nucleotides. The oligonucleotides of the present invention are typically deoxynucleotides. In a preferred embodiment the internucleotide in the oligonucleotide is phosphorodithioate, or more preferably a phosphorothioate bond, although phosphodiester and other internucleotide bonds are within the scope of the invention including oligonucleotides with mixed internucleotide linkages.

Preferred oligonucleotides have the following sequences: The sequences preferably contain all phosphorothioate modified internucleotide linkages.

OLIGO 1: TCC ATG ACG TTC CTG ACG TT

OLIGO 2: TCT CCC AGC GTG CGC CAT

OLIGO 3: ACC GAT GAC GTC GCC GGT GAC GGC ACC ACG

The CpG oligonucleotides utilised in the present invention may be synthesized by any method known in the art (eg EP 468520). Conveniently, such oligonucleotides may be synthesized utilising an automated synthesizer. Methods for producing

phosphorothioate oligonucleotides or phosphorodithioate are described in US5,666,153, US5,278,302 and WO95/26204.

The invention will be further described by reference to the following examples:

5 **EXAMPLE I: Construction of an E. coli strain expressing fusion Protein-D1/3 - E7 -His (HPV16)**

**1) - Construction of expression plasmid**

a) - Plasmid **pMG MCS prot D1/3** (= pRIT14589) is a derivative of pMG81 (described in UK patent application n° 951 3261.9 published as WO97/01640) in  
10 which the codons 4-81 of NS1 coding region from Influenza were replaced by the codons corresponding to residues Ser 20 → Thr127 of mature protein D of Haemophilus Influenzae strain 772, biotype 2 (H. Janson *et al.*, 1991. Infection and Immunity, Jan. p.119-125). The sequence of Prot-D1/3 is followed by a multiple cloning site (11 residues) and a coding region for a C-terminal histidine tail (6 His).

15 This plasmid is used to express the fusion protein D1/3-E7-His.

b) - HPV genomic **E6 and E7 sequences** type HPV 16 (See Dorf *et al.*, Virology 1985, 145, p. 181-185) were amplified from HPV 16 full length genome cloned in pBR322 (obtained from Deutsches Krebsforschungszentrum (DKFZ),  
Referenzzentrum für human pathogen Papillomaviruses - D 69120 - Heidelberg) and  
20 were subcloned into pUC19 to give TCA 301 (= pRIT14462).

**Construction of plasmid TCA 308 (= pRIT14501): a plasmid expressing the fusion Protein-D1/3-E7-His**

The nucleotides sequences corresponding to amino acids 1 → 98 of E7 protein are amplified from pRIT14462. During the polymerase chain reaction, NcoI  
25 and SpeI restriction sites were generated at the 5' and 3' ends of the E7 sequences allowing insertion into the same sites of plasmid pMG MCS Prot D1/3 to give plasmid TCA308 (= pRIT14501). The insert was sequenced to verify that no modification had been generated during the polymerase chain reaction. The sequence for the fusion protein-D1/3-E7-His (HPV 16) is described in sequence ID No.1 and the coding  
30 sequence in ID No.2.

**2) - Transformation of AR58 strain**

Plasmid pRIT14501 was introduced into *E. coli* AR58 (Mott *et al.*, 1985, Proc. Natl. Acad. Sci., 82:88) a defective  $\lambda$  lysogen containing a thermosensitive repressor of the  $\lambda$  pL promoter.

### 3) - Growth and induction of bacterial strain - Expression of Prot -D1/3-E7-His

5 Cells of AR58 transformed with plasmid pRIT14501 were grown in 100 ml of LB medium supplemented with 50  $\mu$ g/ml of Kanamycin at 30°C. During the logarithmic phase of growth bacteria were shifted to 39°C to inactivate the  $\lambda$  repressor and turn on the synthesis of protein D1/3-E7-His. The incubation at 39°C was continued for 4 hours. Bacteria were pelleted and stored at -20°C.

## 10 EXAMPLE II: Construction of an *E.coli* strain expressing fusion Protein-D1/3-E6-his / HPV16

### 1. Construction of expression plasmid

a) Plasmid pMG MCS prot D1/3 (= pRIT14589) is a derivative of pMG81 (described in WO97/01640 in which the codons 4-81 of NS1 coding region from Influenza were replaced by the codons corresponding to residues Ser 20  $\rightarrow$  Thr 127 of mature protein D of Haemophilus Influenzae strain 772, biotype 2 (H. Janson *et al.*, 1991, Infection and Immunity, Jan. p.119-125). The sequence of Prot-D1/3 is followed by a multiple cloning site (11 residues) and a coding region for a C-terminal histidine tail (6 His). This plasmid is used to express the fusion protein D1/3-E6-his.

20 b) HPV genomic E6 and E7 sequences type HPV16 (Seedorf *et al.*, Virology 1985, 145, p.181-185) were amplified from HPV16 full length genome cloned in pBR322 (obtained from Deutsches Krebsforschungszentrum (DKFZ), Referenzzentrum für human pathogen Papillomaviruses -

c) D 69120 - Heidelberg) and were subcloned into pUC19 to give TCA 301 (= pRIT14462).

### 25 Construction of plasmid TCA 307 (=pRIT14497) : a plasmid expressing the fusion Protein-D1/3-E6-His /HPV16

The nucleotides sequences corresponding to amino acid.

1  $\rightarrow$  151 of E6 protein were amplified from pRIT14462. During the polymerase chain reaction, NcoI and SpeI restriction sites were generated at the 5' and 3' ends of the E6 sequences allowing insertion into the same sites of plasmid pMGMCs Prot D1/3 to give plasmid TCA307 (= pRIT14497). The insert was

sequenced to verify that no modification had been generated during the polymerase chain reaction. The protein and coding sequence for the fusion protein-D1/3-E6-His is described in sequence ID No.3 and 4.

## 2. Transformation of AR58 strain

5 Plasmid pRIT14497 was introduced into *E. coli* AR58 (Mott et al., 1985, Proc. Natl. Acad. Sci., 82:88) a defective  $\lambda$  lysogen containing a thermosensitive repressor of the  $\lambda$  pL promoter.

## 3. Growth and induction of bacterial strain - Expression of Prot-D1/3-E6-His

10 Cells of AR58 transformed with plasmid pRIT14497 were grown in 100 ml of LB medium supplemented with 50  $\mu$ g/ml of Kanamycin at 30°C. During the logarithmic phase of growth bacteria were shifted to 39°C to inactivate the  $\lambda$  repressor and turn on the synthesis of protein D1/3-E6-his. The incubation at 39°C was continued for 4 hours. Bacteria were pelleted and stored at -20°C.

## 4. Characterization of fusion Protein D1/3-E6-his (HPV 16)

### 15 Preparation of extracts

Frozen cells are thawed and resuspended in 10 ml of PBS buffer. Cells are broken in a French pressure cell press SLM Aminco at 20.000 psi (three passages). The extract is centrifuged at 16.000 g for 30 minutes at 4°C.

### Analysis on Coomassie-stained SDS-polyacrylamide gels and Western blots

20 After centrifugation of extracts described above, aliquots of supernatant and pellet were analysed by SDS-polyacrylamide gel electrophoresis and Western blotting.

A major band of about 32 kDa, localized in the pellet fraction, was visualised by Coomassie stained gels and identified in Western blots by rabbit polyclonal anti-protein-D and by Ni-NTA conjugate coupled to calf intestinal alkaline phosphatase (Qiagen cat. n° 34510) which detects accessible histidine tail. The level of expression represents about 5 % of total protein.

## 5. Coexpression with thioredoxin

30 In an analogous fashion to the expression of prot D 1/3 E7 His from HPV 18 (example IX) an *E. coli* strain AR58 was transformed with a plasmid encoding thioredoxin and protein D 1/3 E7 His (HPV 16).

**EXAMPLE III: Construction of an *E. coli* strain expressing fusion Protein-D1/3-E6E7-his / HPV16**

**1. Construction of expression plasmid**

- a) Plasmid pMG MCS prot D1/3 (= pRIT14589) is a derivative of pMG81 (described  
5 Supra) in which the codons 4-81 of NS1 coding region from Influenza were replaced  
by the codons corresponding to residues Ser 20 → Thr 127 of mature protein D of  
Haemophilus Influenzae strain 772, biotype 2 (H. Janson *et al.*, 1991, Infection and  
Immunity, Jan. p.119-125). The sequence of Prot-D1/3 is followed by a multiple  
cloning site (11 residues) and a coding region for a C-terminal histidine tail (6 His).  
10 This plasmid is used to express the fusion protein D1/3-E6E7-his.  
b) HPV genomic E6 and E7 sequences type HPV16 (Seedorf *et al.*, Virology 1985,  
145, p.181-185) were amplified from HPV16 full length genome cloned in pBR322  
(obtained from Deutsches Krebsforschungszentrum (DKFZ), Referenzzentrum für  
human pathogen Papillomaviruses - D 69120 - Heidelberg) and were subcloned into  
15 pUC19 to give TCA 301 (= pRIT14462).  
c) The coding sequences for E6 and E7 in TCA301 (= pRIT  
14462) were modified with a synthetic oligonucleotides adaptor (inserted between Afl  
III and Nsi I sites) introducing a deletion of 5 nucleotides between E6 and E7 genes  
to remove the stop codon of E6 and create fused E6 and E7 coding sequences in the  
20 plasmid TCA309(= pRIT 14556).

**Construction of plasmid TCA 311(= pRIT14512) : a plasmid expressing the  
fusion Protein-D1/3-E6E7-His /HPV16**

- The nucleotides sequences corresponding to amino acids 1 → 249 of fused  
E6E7 protein were amplified from pRIT14556. During the polymerase chain  
25 reaction, NcoI and SpeI restriction sites were generated at the 5' and 3' ends of the  
E6E7 fused sequences allowing insertion into the same sites of plasmid pMG MCS  
Prot D1/3 to give plasmid TCA311 (= pRIT14512). The insert was sequenced to  
verify that no modification had been generated during the polymerase chain reaction.  
The protein and coding sequence for the fusion protein-D E6/E7 1/3-His is described  
30 sequence ID No. 5 and 6.

**2. Transformation of AR58 strain**

Plasmid pRIT14512 was introduced into *E. coli* AR58 (Mott et al., 1985, Proc. Natl. Acad. Sci., 82:88) a defective  $\lambda$  lysogen containing a thermosensitive repressor of the  $\lambda$  pL promoter.

### 3. Growth and induction of bacterial strain - Expression of Prot-D1/3-E6E7-His

5 Cells of AR58 transformed with plasmid pRIT14512 were grown in 100 ml of LB medium supplemented with 50  $\mu$ g/ml of Kanamycin at 30°C. During the logarithmic phase of growth bacteria were shifted to 39°C to inactivate the  $\lambda$  repressor and turn on the synthesis of protein D1/3-E6E7-his. The incubation at 39°C was continued for 4 hours. Bacteria were pelleted and stored at -20C.

### 10 4. Characterization of fusion Protein D1/3-E6E7-his

Frozen cells are thawed and resuspended in 10 ml of PBS buffer. Cells are broken in a French pressure cell press SLM Aminco at 20.000 psi (three passages). The extract is centrifuged at 16.000 g for 30 minutes at 4°C.

15 After centrifugation of extracts described above, aliquots of supernatant and pellet were analysed by SDS-polyacrylamide gel electrophoresis and Western blotting.

A major band of about 48 kDa, localized in the pellet fraction, was visualised by Coomassie stained gels and identified in Western blots by rabbit polyclonal anti-protein-D and by Ni-NTA conjugate coupled to calf intestinal alkaline phosphatase  
20 (Qiagen cat. n° 34510) which detects accessible histidine tail. The level of expression represents about 1 % of total protein.

### EXAMPLE: IV

In an analogous fashion the fusion protein of Lipo D 1/3 and E6-E7 from HPV16 was expressed in *E. coli* in the presence of thioredoxin.

25 The N-terminal of the pre-protein (388 aa) contains MDP residues followed by 16 amino acids of signal peptide of lipoprotein D (from Haemophilus Influenzae) which is cleaved in vivo to give the mature protein (370 aa). Lipoprotein portion (aa 1 to 127) is followed by the proteins E6 and E7 in fusion. The C terminal of the protein is elongated by TSGHHHHHH.

### 30 EXAMPLE V: Construction of E.coli strain B1002 expressing fusion ProtD1/3-E7

Mutated (cys24->gly,glu26->gln ) type HPV16



# 1)-Construction of expression plasmid

## Starting material:

- a) - Plasmid pRIT 14501 ( = TCA 308) which codes for fusion ProtD1/3-E7 -His
- b) - Plasmid LITMUS 28 ( New England Biolabs cat n° 306-28 ), a cloning vector
- 5 pUC-derived
- c) - Plasmid pMG MCS ProtD1/3 ( pRIT 14589) , a derivative of pMG81 (described Supra) in which the codons 4-81 of NS1 coding region from Influenza were replaced by the codons corresponding to residues Ser 20 → Thr 127 of mature protein D of Haemophilus Influenzae strain 772, biotype 2 (H. Janson *et al.*, 1991, Infection and
- 10 Immunity, Jan. p.119-125). The sequence of Prot-D1/3 is followed by a multiple cloning site (11 residues) and a coding region for a C-terminal histidine tail (6 His)

## Construction of plasmid pRIT 14733( =TCA347): a plasmid expressing the fusion Protein-D1/3-E7 mutated ( cys24->gly ,glu26->gln) with His tail

- The NcoI - XbaI fragment from pRIT 14501 (=TCA 308 ), bearing the coding
- 15 sequence of E7 gene from HPV16 , elongated with an His tail , was subcloned in an intermediate vector Litmus 28 useful for mutagenesis to give pRIT 14909 (=TCA337)
  - Double mutations cys24->gly ( Edmonds and Vousden , J.Virology 63 : 2650 (1989) and glu26->gln ( Phelps et al , J.Virology 66: 2418-27 ( 1992 ) were chosen to impair the binding to the antioncogene product of Retinoblastome gene ( pRB ).
  - 20 The introduction of mutations in E7 gene was realized with the kit " Quick Change Site directed Mutagenesis ( Stratagene cat n° 200518) to give plasmid pRIT 14681(=TCA343 ) .After verification of presence of mutations and integrity of the complete E7 gene by sequencing , the mutated E7 gene was introduced into vector pRIT 14589 ( = pMG MCS ProtD1/3 ) to give plasmid pRIT 14733 (=TCA347)
  - 25 protein and coding sequence.

The sequence for the fusion protein-D1/3-E 7 mutated ( cys24->gly, glu26->gln ) -His is described in sequence ID No. 7 and 8.

## 2)-Construction of strain B1002 expressing ProtD1/3-E7mutated (cys 24->gly , glu26->gln )-His /HPV16

- 30 Plasmid pRIT 14733 was introduced into *E.coli* AR58 ( Mott et al. ,1985,

Proc. Natl. Acad. Sci. , 82:88 ) a defective  $\lambda$  lysogen containing a thermosensitive repressor of the  $\lambda$  pL promoter ,to give strain B1002 , by selection for transformants resistant to kanamycine

### 3)-Growth and induction of bacterial strain B1002 - Expression of ProtD1/3-E7

#### 5 mutated (cys 24->gly , glu26->gln )-His /HPV16

Cells of AR58 transformed with plasmid pRIT 14733 ( B1002 strain ) were grown at 30°C in 100 ml of LB medium supplemented with 50  $\mu$ gr /ml of Kanamycin. During the logarithmic phase of growth bacteria were shifted to 39°C to inactivate the  $\lambda$  repressor and turn on the synthesis of ProtD1/3-E7 mutated -His /HPV16 . The  
10 incubation at 39°C was continued for 4 hours . Bacteria were pelleted and stored at -20°C.

#### 4)-Characterization of fusion ProtD1/3-E7 mut (cys24->gly, glu26->gln)- His type HPV16.

Frozen cells were thawed and resuspended in 10 ml of PBS buffer. Cells were  
15 broken in a French Pressure cell press SLM Aminco at 20 000 psi ( three passages) . The extract was centrifuged at 16000 g for 30 minutes at 4°C.

After centrifugation of extracts described above, aliquots of supernatant and pellet were analysed by SDS-polyacrylamide gel electrophoresis and Western blotting.

20 A major band of about 33 kDa, localized in the pellet fraction, was visualised by Coomassie stained gels and identified in Western blots by rabbit polyclonal 22 J 70 anti-protein D, by monoclonal anti E7 /HPV16 from Zymed and by Ni-NTA conjugate coupled to calf intestinal alkaline phosphatase (Qiagen cat. n° 34510) which detects accessible histidine tail. The level of expression represents about 3 to 5 % of  
25 total protein.

Cells of B1002 were separated from the culture broth by centrifugation.

The concentrated cells of B1002 were stored at -65°C.

### EXAMPLE VI: Construction of an *E. coli* strain expressing fusion clyta-E6-his (HPV 16)

#### 30 1. Construction of expression plasmid

a) -Plasmid pRIT14497 (= TCA307), that codes for fusion ProtD1/3-E6-His /HPV16

b)-Plasmid pRIT14661 (= DVA2), an intermediate vector containing the coding sequence for the 117 C-terminal codons of LytA of *Streptococcus Pneumoniae*. LytA is derived from *Streptococcus pneumoniae* which synthesize an N-acetyl-L-alanine amidase, amidase LYTA. (coded by the *lytA* gene {Gene, 43 (1986) pag 265-272} an autolysin that specifically degrades certain bonds in the peptidoglycan backbone . The C-terminal domain of the LYTA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE.

**1.b Construction of plasmid pRIT14634 (=TCA332): a plasmid expressing the fusion clyta-E6-His /HPV16**

10 a)The first step was the purification of the large NcoI-AflIII restriction fragment from plasmid pRIT14497 and the purification of the small AflII-AflIII restriction fragment from pRIT14661

b)The second step was linking of clyta sequences to the E7-His sequences (NcoI and AflIII are compatible restriction sites) that gave rise to the plasmid pRIT 14634  
15 (=TCA332), coding for the fusion protein clyta-E6-His under the control of the pL promoter.

The protein and coding sequence for the fusion protein clyta-E6-His is described sequence ID No. 9 and 10.

**Transformation of AR58 strain**

20 Plasmid pRIT14634 was introduced into *E. coli* AR58 (Mott et al., 1985, Proc. Natl. Acad. Sci., 82:88) a defective  $\lambda$  lysogen containing a thermosensitive repressor of the  $\lambda$  pL promoter.

**Growth and induction of bacterial strain - Expression of clyta-E6-His**

Cells of AR58 transformed with plasmid pRIT14634 were grown in 100 ml of  
25 LB medium supplemented with 50  $\mu$ g/ml of Kanamycin at 30°C. During the logarithmic phase of growth bacteria were shifted to 39°C to inactivate the  $\lambda$  repressor and turn on the synthesis of protein clyta-E6-his. The incubation at 39°C was continued for 4 hours. Bacteria were pelleted and stored at -20°C.

**4. Characterization of fusion clyta-E6-his**

30 Frozen cells were thawed and resuspended in 10 ml of PBS buffer. Cells were broken in a French pressure cell press SLM Aminco at 20.000 psi (three passages). The extract was centrifuged at 16.000 g for 30 minutes at 4°C. After centrifugation of

extracts described above, aliquots of supernatant and pellet were analysed by SDS-polyacrylamide gel electrophoresis and Western blotting.

A major band of about 33 kDa, localized in the pellet fraction, was visualised by Coomassie stained gels and identified in Western blots by rabbit polyclonal anti-clyta antibodies and by Ni-NTA conjugate coupled to calf intestinal alkaline phosphatase (Qiagen cat. n° 34510) which detects accessible histidine tail. The level of expression represents about 3 % of total protein.

**EXAMPLE VII: Construction of an *E. coli* strain expressing fusion clyta-E7-his (HPV 16)**

**1. Construction of expression plasmid**

**1.a Starting materials**

- a) -Plasmid pRIT14501 (= TCA308), that codes for fusion ProtD1/3-E7-His /HPV16
- b)-Plasmid pRIT14661 (= DVA2), an intermediate vector containing the coding sequence for the 117 C-terminal codons of LytA of *Streptococcus Pneumoniae*.

**1.b Construction of plasmid pRIT14626 (=TCA330): a plasmid expressing the fusion clyta-E7-His / HPV16**

- a) The first step was the purification of the large NcoI-AflIII restriction fragment from plasmid pRIT14501 and the purification of the small AflIII-AflIII restriction fragment from pRIT14661
- b) The second step was linking of clyta sequences to the E7-His sequences (NcoI and AflIII are compatible restriction sites) that gave rise to the plasmid pRIT 14626 (=TCA330), coding for the fusion protein clyta-E7-His under the control of the pL promoter.

The protein and coding sequence for the fusion protein clyta-E7-His is described in sequence ID No. 11 and 12.

**2. Transformation of AR58 strain**

Plasmid pRIT14626 was introduced into *E. coli* AR58 (Mott et al., 1985, Proc. Natl. Acad. Sci., 82:88) a defective  $\lambda$  lysogen containing a thermosensitive repressor of the  $\lambda$  pL promoter.

**3. Growth and induction of bacterial strain - Expression of clyta-E7-His**

Cells of AR58 transformed with plasmid pRIT14626 were grown in 100 ml of LB medium supplemented with 50  $\mu$ gr/ml of Kanamycin at 30°C. During the

logarithmic phase of growth bacteria were shifted to 39°C to inactivate the  $\lambda$  repressor and turn on the synthesis of protein clyta-E7-his. The incubation at 39°C was continued for 4 hours. Bacteria were pelleted and stored at -20°C.

#### 4. Characterization of fusion clyta-E7-his

5 Frozen cells were thawed and resuspended in 10 ml of PBS buffer. Cells were broken in a French pressure cell press SLM Aminco at 20.000 psi (three passages). The extract was centrifuged at 16.000 g for 30 minutes at 4°C. After centrifugation of extracts described above, aliquots of supernatant and pellet were analysed by SDS-polyacrylamide gel electrophoresis and Western blotting.

10 A major band of about 35 kDa, localized in the pellet fraction, was visualised by Coomassie stained gels and identified in Western blots by rabbit polyclonal anti-clyta antibodies and by Ni-NTA conjugate coupled to calf intestinal alkaline phosphatase (Qiagen cat. n° 34510) which detects accessible histidine tail. The level of expression represents about 5 % of total protein.

#### 15 EXAMPLE VIII: Construction of an *E. coli* strain expressing fusion clyta-E6E7-his (HPV 16)

##### 1. Construction of expression plasmid

##### 1.a Starting materials

- 20 a) -Plasmid pRIT14512 (= TCA311), that codes for fusion ProtD1/3-E6E7-His /HPV16
- b)-Plasmid pRIT14661 (= DVA2), an intermediate vector containing the coding sequence for the 117 C-terminal codons of LytA of *Streptococcus Pneumoniae*.

##### 1.b Construction of plasmid pRIT14629 (=TCA331): a plasmid expressing the fusion clyta-E6E7-His /HPV16

- 25 a)The first step was the purification of the large NcoI-AflIII restriction fragment from plasmid pRIT14512 and the purification of the small AflII-AflIII restriction fragment from pRIT14661
- b)The second step was linking of clyta sequences to the E7-His sequences (NcoI and AflIII are compatible restriction sites)that gave rise to the plasmid pRIT 14629
- 30 (=TCA331), coding for the fusion protein clyta-E6E7-His under the control of the pL promoter.

The protein and coding sequence for the fusion protein clyta-E6E7-His is sequenced ID No. 13 and 14.

## 2. Transformation of AR58 strain

Plasmid pRIT14629 was introduced into *E. coli* AR58 (Mott et al., 1985, Proc. Natl. Acad. Sci., 82:88) a defective  $\lambda$  lysogen containing a thermosensitive repressor of the  $\lambda$  pL promoter.

## 3. Growth and induction of bacterial strain - Expression of clyta-E6E7-His

Cells of AR58 transformed with plasmid pRIT14629 were grown in 100 ml of LB medium supplemented with 50  $\mu$ g/ml of Kanamycin at 30°C. During the logarithmic phase of growth bacteria were shifted to 39°C to inactivate the  $\lambda$  repressor and turn on the synthesis of protein clyta-E6E7-his. The incubation at 39°C was continued for 4 hours. Bacteria were pelleted and stored at -20°C.

## 4. Characterization of fusion clyta-E6E7-his

Frozen cells were thawed and resuspended in 10 ml of PBS buffer. Cells were broken in a French pressure cell press SLM Aminco at 20.000 psi (three passages). The extract was centrifuged at 16.000 g for 30 minutes at 4°C.

After centrifugation of extracts described above, aliquots of supernatant and pellet were analysed by SDS-polyacrylamide gel electrophoresis and Western blotting.

A major band of about 48 kDa, localized in the pellet fraction, was visualised by Coomassie stained gels and identified in Western blots by rabbit polyclonal anti-clyta antibodies and by Ni-NTA conjugate coupled to calf intestinal alkaline phosphatase (Qiagen cat. n° 34510) which detects accessible histidine tail. The level of expression represents about 1 % of total protein.

## EXAMPLE IX: Prot D1/3 E7 his (HPV 18) (E.Coli B1011)

Protein D1/3 E7 his HPV expressed with Thioredoxin inTrans (E.Coli B1012)

### 1) - Construction of expression plasmids

1).a.Construction of plasmid TCA316(=pRIT 14532) a plasmid expressing the fusion Protein-D1/3-E7-His /HPV18

### 30 Starting materials

a) - Plasmid pMG MCS prot D1/3 (= pRIT14589) is a derivative of pMG81 (described in UK patent application n° 951 3261.9 published as WO97/01640 in

which the codons 4-81 of NS1 coding region from Influenza were replaced by the codons corresponding to residues Ser 20 → Thr 127 of mature protein D of Haemophilus Influenzae strain 772, biotype 2 (H. Janson *et al.*, 1991, Infection and Immunity, Jan. p.119-125). The sequence of Prot-D1/3 is followed by a multiple cloning site (11 residues) and a coding region for a C-terminal histidine tail (6 His).  
 5 This plasmid is used to express the fusion protein D1/3-E7-his.

b) - HPV genomic E6 and E7 sequences of prototype HPV18(Cole *et al.*J.Mol.Biol.(1987)193,599-608) were amplified from HPV16 full length genome cloned in pBR322 (obtained from Deutsche Krebsforschungszentrum (DKFZ),  
 10 Referenzzentrum für human pathogen Papillomaviruses - D 69120 - Heidelberg) and were subcloned into pUC19 to give TCA 302 (= pRIT14467).

#### Construction of plasmid TCA 316(= pRIT14532)

The nucleotides sequences corresponding to amino acids 1 → 105 of E7 protein were amplified from pRIT14467. During the polymerase chain reaction, NcoI and SpeI restriction sites were generated at the 5' and 3' ends of the E7 sequences  
 15 allowing insertion into the same sites of plasmid pMGMCs Prot D1/3 to give plasmid TCA316 (= pRIT14532). The insert was sequenced and a modification versus E7/HPV18 prototype sequence was identified in E7 gene (nucleotide 128 G→A) generating a substitution of a glycine by a glutamic acid (aa 43 in E7, position 156 in  
 20 fusion protein). The protein and coding sequence for the fusion protein-D1/3-E7-His /HPV18 is set forth in sequence ID No. 15 and ID No. 16.

#### 1).b. Construction of plasmid TCA313 (=pRIT14523): a plasmid expressing thioredoxin

##### Starting materials

- 25 a) - Plasmid pBBR1MCS4(Antoine R. and C.Locht,Mol.Microbiol. 1992,6,1785-1799 ; M.E.Kovach *et al.* Biotechniques 16, (5), 800-802 )which is compatible with plasmids containing ColE1 or P15a origins of replication.
- b) - Plasmid pMG42 (described in WO93/04175) containing the sequence of promoter pL of Lambda phage
- 30 c) - Plasmid pTRX ( Invitrogen, kit Thiofusion K350-01) bearing the coding sequence for thioredoxin followed by AspA transcription terminator.

#### Construction of plasmid TCA313(=pRIT14523)

The fragment EcoRI-NdeI fragment from pMG42, bearing pL promoter and the NdeI-HindIII fragment from pTRX, bearing the coding sequence for thioredoxin followed by AspA terminator, were purified and ligated into the EcoRI and HindIII sites of plasmid vector pBBR1MCS4 to give plasmid TCA313(= pRIT14523).

5 The coding sequence for thioredoxin is described in ID No. 17.

## 2) - Transformation of AR58 strain

### 2).a. To obtain strain B1011 expressing ProtD1/3-E7-His/HPV18

Plasmid pRIT14532 was introduced into *E. coli* AR58 (Mott et al., 1985. Proc. Natl. Acad. Sci., 82:88) a defective  $\lambda$  lysogen containing a thermosensitive repressor of the  
10  $\lambda$  pL promoter, by selection for transformants resistant to kanamycine.

### 2).b. Construction of strain B1012 expressing ProtD1/3-E7-His/HPV18 and thioredoxin

Plasmid pRIT14532 and pRIT14523 were introduced into *E. coli* AR58 (Mott et al., 1985, Proc. Natl. Acad. Sci., 82:88) a defective  $\lambda$  lysogen containing a  
15 thermosensitive repressor of the  $\lambda$  pL promoter, by double selection for transformants resistant to kanamycin and ampicillin.

## 3) - Growth and induction of bacterial strains B1011 and B1012 - Expression of Prot-D1/3-E7-His/HPV18 without and with thioredoxin in trans

Cells of AR58 transformed with plasmids pRIT14532 (B1011 strain) and  
20 Cells of AR58 transformed with plasmids pRIT14532 and pRIT14523 (B1012 strain) were grown at 30°C in 100 ml of LB medium supplemented with 50  $\mu$ g/ml of Kanamycin for B1011 strain and supplemented 50  $\mu$ g/ml of Kanamycin and 100  $\mu$ g/ml of Ampicillin for B1012 strain. During the logarithmic phase of growth bacteria were shifted to 39°C to inactivate the  $\lambda$  repressor and turn on the synthesis  
25 of protein D1/3-E7-his/HPV18 and thioredoxin. The incubation at 39°C was continued for 4 hours.

### Characterization of fusion Protein D1/3-E7-his /HPV18

#### Preparation of extracts

Frozen cells are thawed and resuspended in 10 ml of PBS buffer. Cells are  
30 broken in a French pressure cell press SLM Aminco at 20.000 psi (three passages). The extract is centrifuged at 16.000 g for 30 minutes at 4°C.

Analysis n Co massie-stained SDS-polyacrylamide gels and Western blots



After centrifugation of extracts described above, aliquots of supernatant and pellet were analysed by SDS-polyacrylamide gel electrophoresis and Western blotting.

The fusion protD1/3-E7-His (about 31 kDa) was visualised by Coomassie stained gels in the pellet fraction for strain B1011 and partially localized (30%) in the supernatant fraction for strain B1012 and was identified in Western blots by rabbit polyclonal anti-protein-D and by Ni-NTA conjugate coupled to calf intestinal alkaline phosphatase (Qiagen cat. n° 34510) which detects accessible histidine tail. The level of expression represents about 1-3% of total protein as shown on a Coomassie-stained SDS-polyacrylamide gel.

For the extract of strain B1012 the thioredoxin ( about 12 KDa) was visualised by coomassie stained gel in the supernatant and identified in western blots by monoclonal anti thioredoxin ( Invitrogen R920-25)

**EXAMPLE X: Construction of E.coli strain B1098 expressing fusion ProtD1/3-**

**E7**

**Mutated (cys27->gly,glu29->gln ) type HPV18**

**1)-Construction of expression plasmid**

**Starting material:**

- a) - Plasmid pRIT 14532 ( = TCA 316) which codes for fusion ProtD1/3-E7 -His
- b) - Plasmid LITMUS 28 ( New England Biolabs cat n° 306-28 ) , a cloning vector pUC-derived
- c) - Plasmid pMG MCS ProtD1/3 ( pRIT 14589) , a derivative of pMG81 (described supra) in which the codons 4-81 of NS1 coding region from Influenza were replaced by the codons corresponding to residues Ser 20 → Thr 127 of mature protein D of Haemophilus Influenzae strain 772, biotype 2 (H. Janson *et al.*, 1991, Infection and Immunity, Jan. p.119-125). The sequence of Prot-D1/3 is followed by a multiple cloning site (11 residues) and a coding region for a C-terminal histidine tail (6 His)

**Construction of plasmid pRIT 14831( =TCA355): a plasmid expressing the fusion Protein-D1/3-E7 mutated ( cys27->gly ,glu29->gln) with His tail**

The NcoI - XbaI fragment from pRIT 14532 (=TCA 316 ), bearing the coding sequence of E7 gene from HPV18 , elongated with an His tail , was subcloned in an intermediate vector Litmus 28 useful for mutagenesis to give pRIT 14910 (=TCA348)

By analogy with E7/HPV16 mutagenesis, double mutations *cys27-->gly* and *glu29-->gln* were chosen to impair the binding to the antioncogene product of Retinoblastome gene ( pRB ).

The introduction of mutations in E7 gene was realized with the kit " Quick Change Site directed Mutagenesis ( Stratagene cat n° 200518) .As the sequencing of pRIT14532 had pointed out the presence of a glutamic acid in position 43 of E7 instead of a glycine in the prototype sequence of HPV18 , a second cycle of mutagenesis was realized to introduce a glycine in position 43 . We obtained plasmid pRIT 14829 ( = TCA353 ). After verification of presence of mutations and integrity of the complete E7 gene by sequencing , the mutated E7 gene was introduced into vector pRIT 14589 ( = pMG MCS ProtD1/3 ) to give plasmid pRIT 14831 (=TCA355).

The protein and coding sequence for the fusion protein-D1/3-E 7 mutated (*cys27->gly*, *glu29->gln* ) -His is described in sequence ID No. 18 and 19.

2)Construction of strain B1098 expressing ProtD1/3-E7mutated (*cys 27->gly* , *glu29->gln* )-His /HPV18

Plasmid pRIT 14831 was introduced into *E.coli* AR58 ( Mott et al. ,1985, Proc. Natl. Acad. Sci. , 82:88 ) a defective  $\lambda$  lysogen containing a thermosensitive repressor of the  $\lambda$  pL promoter ,to give strain B1098 , by selection for transformants resistant to kanamycin.

3)-Growth and induction of bacterial strain B1098 - Expression of ProtD1/3-E7 mutated (*cys 27->gly* , *glu29->gln* )-His /HPV18

Cells of AR58 transformed with plasmid pRIT 14831 ( B1098 strain ) were grown at 30°C in 100 ml of LB medium supplemented with 50  $\mu$ gr /ml of Kanamycin. During the logarithmic phase of growth bacteria were shifted to 39°C to inactivate the  $\lambda$  repressor and turn on the synthesis of ProtD1/3-E7 mutated -His /HPV18 . The incubation at 39°C was continued for 4 hours. Bacteria were pelleted and stored at - 20°C.

4)-Characterization of fusion ProtD1/3-E7 mut (*cys24->gly*, *glu26->gln*)- His type HPV16

Frozen cells were thawed and resuspended in 10 ml of PBS buffer. Cells were broken in a French Pressure cell press SLM Aminco at 20 000 psi ( three passages) . The extract was centrifuged at 16000 g for 30 minutes at 4°C.

#### Analysis on Coomassie stained SDS-polyacrylamide gels and Western blots

- 5 After centrifugation of extracts described above, aliquots of supernatant and pellet were analysed by SDS-polyacrylamide gel electrophoresis and Western blotting. A major band of about 31 kDa, localized in the pellet fraction, was visualised by Coomassie stained gels and identified in Western blots by rabbit polyclonal 22 J 70 anti-protein D and by monoclonal Penta-His (Qiagen cat. n° 34660) which detects  
10 accessible histidine tail. The level of expression represents about 3 to 5 % of total protein.

#### EXAMPLE XI: Construction of an *E. coli* strain expressing fusion Protein-D1/3-E6-his / HPV18

##### 1. Construction of expression plasmid

- 15 a) Plasmid pMG MCS prot D1/3 (= pRIT14589) is a derivative of pMG81 (described supra) in which the codons 4-81 of NS1 coding region from Influenza were replaced by the codons corresponding to residues Ser 20 → Thr 127 of mature protein D of Haemophilus Influenzae strain 772, biotype 2 (H. Janson *et al.*, 1991, Infection and Immunity, Jan. p.119-125). The sequence of Prot-D1/3 is followed by a multiple  
20 cloning site (11 residues) and a coding region for a C-terminal histidine tail (6 His). This plasmid is used to express the fusion protein D1/3-E6-his. HPV genomic E6 and E7 sequences type HPV18 (Cole et al., J. Mol. Biol. 1987, 193 , p.599-608. ) were amplified from HPV18 full length genome cloned in pBR322 (obtained from Deutsches Krebsforschungszentrum (DKFZ), Referenzzentrum für  
25 human pathogen Papillomaviruses - D 69120 - Heidelberg) and were subcloned into pUC19 to give TCA 302 (= pRIT14467).

##### Construction of plasmid TCA 314(= pRIT14526) : a plasmid expressing the fusion Protein-D1/3-E6-His /HPV18

- The nucleotides sequences corresponding to amino acids  
30 1 → 158 of E6 protein were amplified from pRIT14467. During the polymerase chain reaction, NcoI and SpeI restriction sites were generated at the 5' and 3' ends of the E6 sequences allowing insertion into the same sites of plasmid

pMG MCS Prot D1/3 to give plasmid TCA314 (= pRIT14526). The insert was sequenced to verify that no modification had been generated during the polymerase chain reaction. The protein and coding sequence for the fusion protein-D1/3-E6-His is described in sequence ID No. 20 and 21.

5 **Transformation of AR58 strain**

Plasmid pRIT14526 was introduced into *E. coli* AR58 (Mott et al., 1985, Proc. Natl. Acad. Sci., 82:88) a defective  $\lambda$  lysogen containing a thermosensitive repressor of the  $\lambda$  pL promoter.

**3. Growth and induction of bacterial strain - Expression of Prot-D1/3-E6-His**

10 Cells of AR58 transformed with plasmid pRIT14526 were grown in 100 ml of LB medium supplemented with 50  $\mu$ g/ml of Kanamycin at 30°C. During the logarithmic phase of growth bacteria were shifted to 39°C to inactivate the  $\lambda$  repressor and turn on the synthesis of protein D1/3-E6-his. The incubation at 39°C was continued for 4 hours. Bacteria were pelleted and stored at -20°C.

15 **4. Characterization of fusion Protein D1/3-E6-his**

Frozen cells are thawed and resuspended in 10 ml of PBS buffer. Cells are broken in a French pressure cell press SLM Aminco at 20.000 psi (three passages). The extract is centrifuged at 16.000 g for 30 minutes at 4°C. After centrifugation of extracts described above, aliquots of supernatant and pellet were analysed by SDS-

20 polyacrylamide gel electrophoresis and Western blotting.

A major band of about 32 kDa, localized in the pellet fraction, was visualised by Coomassie stained gels and identified in Western blots by rabbit polyclonal anti-protein-D and by Ni-NTA conjugate coupled to calf intestinal alkaline phosphatase (Qiagen cat. n° 34510) which detects accessible histidine tail. The level of expression

25 represents about 3-5 % of total protein.

**EXAMPLE XII: Construction of an *E. coli* strain expressing fusion Protein-D1/3-E6E7-his / HPV18**

**I. Construction of expression plasmid**

a) Plasmid pMG MCS prot D1/3 (= pRIT14589) is a derivative of pMG81 (described supra) in which the codons 4-81 of NS1 coding region from Influenza were replaced by the codons corresponding to residues Ser 20  $\rightarrow$  Thr 127 of mature protein D of Haemophilus Influenzae strain 772, biotype 2 (H. Janson et al., 1991, Infection and

Immunity, Jan. p.119-125). The sequence of Prot-D1/3 is followed by a multiple cloning site (11 residues) and a coding region for a C-terminal histidine tail (6 His).

This plasmid is used to express the fusion protein D1/3-E6E7-his.

5 b) HPV genomic E6 and E7 sequences type HPV18 (Cole et al., J.Mol.Biol. 1987, 193, 599-608) were amplified from HPV18 full length genome cloned in pBR322 (obtained from Deutsches Krebsforschungszentrum (DKFZ), Referenzzentrum für human pathogen Papillomaviruses - D 69120 - Heidelberg) and were subcloned into pUC19 to give TCA 302 (= pRIT14467).

10 c) The coding sequences for E6 and E7 in TCA302 (= pRIT 14467) were modified with a synthetic oligonucleotides adaptor (inserted between Hga I and Nsi I sites) introducing a deletion of 11 nucleotides between E6 and E7 genes, removing the stop codon of E6 and creating fused E6 and E7 coding sequences in the plasmid TCA320( = pRIT 14618 ).

15 **Construction of plasmid TCA 328(= pRIT14567) : a plasmid expressing the fusion Protein-D1/3-E6E7-His /HPV18**

The nucleotides sequences corresponding to amino acids

20 1 → 263 of fused E6E7 protein were amplified from pRIT14618. During the polymerase chain reaction, NcoI and SpeI restriction sites were generated at the 5' and 3' ends of the E6E7 fused sequences allowing insertion into the same sites of plasmid pMGMCS Prot D1/3 to give plasmid TCA328 (= pRIT14567). The insert was sequenced to verify that no modification had been generated during the polymerase chain reaction. The protein and coding sequence for the fusion protein-D1/3-E6E7-His is described in sequence ID No. 22 and 23.

## **2. Transformation of AR58 strain**

25 Plasmid pRIT14567 was introduced into *E. coli* AR58 (Mott et al., 1985, Proc. Natl. Acad. Sci., 82:88) a defective  $\lambda$  lysogen containing a thermosensitive repressor of the  $\lambda$  pL promoter.

## **3. Growth and induction of bacterial strain - Expression of Prot-D1/3-E6E7-His**

30 Cells of AR58 transformed with plasmid pRIT14512 were grown in 100 ml of LB medium supplemented with 50  $\mu$ g/ml of Kanamycin at 30°C. During the logarithmic phase of growth bacteria were shifted to 39°C to inactivate the  $\lambda$  repressor

and turn on the synthesis of protein D1/3-E6E7-his. The incubation at 39°C was continued for 4 hours. Bacteria were pelleted and stored at -20°C.

#### 4.Characterization of fusion Protein D1/3-E6E7-his

Frozen cells are thawed and resuspended in 10 ml of PBS buffer. Cells are broken in a French pressure cell press SLM Aminco at 20.000 psi (three passages). The extract is centrifuged at 16.000 g for 30 minutes at 4°C.

After centrifugation of extracts described above, aliquots of supernatant and pellet were analysed by SDS-polyacrylamide gel electrophoresis and Western blotting.

A major band of about 48 kDa, localized in the pellet fraction, was visualised by Coomassie stained gels and identified in Western blots by rabbit polyclonal anti-protein-D and by Ni-NTA conjugate coupled to calf intestinal alkaline phosphatase (Qiagen cat. n° 34510) which detects accessible histidine tail. The level of expression represents about 1 % of total protein.

#### 15 EXAMPLE XIII

The therapeutic potential of vaccine containing the PD1/3 E7 fusion protein and different CpG oligonucleotides were evaluated in the TC1 (E7 expressing tumour model.)

##### 1. Therapeutic experiments: protocol

10e6 TC1 cells, E7 expressing tumour cells: were injected subcutaneously (200µl) in the flank of C57BL/6 immunocompetent mice. Mice were vaccinated 7 and 14 days after the tumour challenge, with 5µg ProtD 1/3 E7 HPV16 injected intra-footpad (100µl : 50µl / footpad) in the presence of different adjuvants:

2 and 4 weeks after the second immunisation, 5 mice/group were killed and spleens or popliteal lymph nodes were taken and analyzed for immune response.

##### 1.2 Results

###### Groups of mice

- 1) PBS
- 2) ProtD1/3 E7 HPV16
- 30 3) ProtD1/3 E7 HPV16 + oligo 1: 1826 (WD 1001): TCC ATG ACG TTC CTG ACG TT
- 4) Oligo 1
- 5) ProtD1/3 E7 HPV16 + oligo 2/ 1758 (WD1002): TCT CCC AGC GTG CGC CAT

## 6) Oligo 2

**Tumour Growth;**

was monitored by measuring individual tumours twice a week.

Figure 1 : represents the mean tumour growth (in mm<sup>2</sup>)/group n=10 followed  
5 over 4 weeks.

- The injection of 10e6 TC1 cells injected subcutaneously give rise to a growing tumour in 100% of the animals.
- Vaccinating with ProtD1/3E7 or adjuvant alone: 100% of the animals develop a tumour.
- 10 • As shown in figure 1 and 2, in the groups of mice that received the antigen with a CpG oligonucleotide the mean tumour growth remained very low and very similar between groups, reflecting that the tumour growth either was slowed down or that several tumours were completely rejected.

The analysis of individual tumour growth 2 and 4 weeks after the latest  
15 vaccination showed that complete rejection in the groups were:

	Day 28 (n=10)	day 42 (n=5)
E7+oligo1 (1826)	40%	40%
Oligo1	0%	0%
E7+oligo2 (1758)	70%	40%
Oligo2	0%	0%

The mean tumour growth/group of mice vaccinated with PD1/3 E7+ the CpG oligos are quite similar and analysis of the individual tumour growth showed that the  
20 CpG oligos induce prolonged complete tumour rejection.

**Conclusion**

Both CpG (Oligo 2>oligo 1) induced complete tumour regression .

Lymphoproliferative response was analysed by *in vitro* restimulation of spleen and lymph nodes cells for 72 hrs with either PD1/3E7, the protein E7(Bollen)  
25 and PD (whole) PD1/3 (coated or not on latex  $\mu$ beads) (10, 1, 0.1  $\mu$ g/ml) 2 and 4 weeks post II.

- Positive controls (ConA stimulation) were positive.

- Surprisingly, no E7 specific and no PD specific proliferative response could be observed starting with spleen cells 2 or 4 weeks post II (probably due to a technical problem: data not shown).
- On the contrary, lymph node cells from mice that received ProtD1/3 E7 in CpG oligos 1 and 2 showed a very good E7 specific proliferative response although almost no PD (whole) specific response could be observed even at the highest concentration of 100µg/ml no PD1/3 specific responses was observed even when coated on latex µbeads.

Similar data were obtained 4 weeks post II.

#### 10 Serology

The anti E7 antibody response: IgG tot and isotypes (IgG1, IgG2a, IgG2b, IgGTot) were measured by ELISA using the E7 protein as coating antigen as described in the Materials and Methods. Figures 3 and 4 show the relative percentage of the different IgG isotypes in the total of IgGs, 2 and 4 weeks post II respectively.

- The Oligos affect only weakly (oligo 2) or not at all (Oligo 1) the weak antibody response observed when PD1/3E7 alone was injected.
- The predominant E7 specific antibody subclass was clearly IgG2b for all the formulation tested (80-90% of the total IgGs).

The same results were obtained 4 weeks post II

#### 20 Isotypic profile of anti E7 responses (post II, pooled sera) exp. 97293

Groups	IgG1	IgG2a	IgG2b	IgGtot
1) PBS	0	0	0	0
2) ProtD1/3 E7 HPV16	1020	0	4130	4740
3) ProtD1/3 E7 HPV16 + oligo 1	170	400	3680	4910
4) Oligo 1	0	0	530	420
5) ProtD1/3 E7 HPV16 + oligo 2	0	590	7560	13690
6) Oligo 2	0	0	0	0

Groups	IgG1	IgG2a	IgG2b	IgGtot
1) PBS	0	0	0	0
2) ProtD1/3 E7 HPV1	240	0	1650	1400
3) ProtD1/3 E7 HPV16 + oligo 1	0	0	1280	1430
4) Oligo 1	0	0	0	0
5) ProtD1/3 E7 HPV16 + oligo 2	0	560	3600	5880
6) Oligo 2	0	0	0	0



**CTL assay:**

A CTL response could be detected when measured 2 weeks after the latest vaccination, when cells were re-stimulated in vitro with irradiated TC1 when TC1 or peptide E7 pulsed EL4, were used as target cells, when mice immunised with PD1/3 E7 + CpG oligo 2> 1 (25-40% specific lysis) and not with oligos alone.

- Lysis was seen on TC1 cells than on peptide E7 pulsed EL4 cells, but this is mostly observed in the groups of mice vaccinated with PD1/3E7 + CpG oligos (2>1). In this experiment other formulations did not induce a CTL.
- Using E7 pulsed EL4 cells, no lysis was observed when mice received the protein or the adjuvant alone.

**1.3 Materials and Methods**

Component	Brand	Batch number	Concentration (mg/ml)	Buffer
ProtD1/3-E7		957/015	0.677	PBS 7.4
oligo CpG 1826	EuroGentec	WD1001	5	H <sub>2</sub> O
oligo CpG	EuroGentec	WD1002	5	H <sub>2</sub> O

**1.3.1 Formulation Process**

All the formulations were prepared on the day of injection.

**Oligo containing formulations**

Formulations containing oligo alone without other adjuvant were prepared by addition of CpG to the diluted PrtD1/3-E7 in PBS pH 7.4.

The adjuvant controls without antigen were prepared by replacing the protein by PBS.

**1.3.2 Mice and Cell lines**

Mice C57Bl/6 (Iffa Credo) 6-8 weeks old mice were used in these experiments.

**Cell lines:** TC1 (obtained from the John Hopkin's University), or EL4 cells were grown in RPMI 1640 (Bio Whittaker) containig 10% FCS and additives: 2mM L-Glutamine, 1% antibiotics (10000U/ml penicilin, 10000µg/ml streptomycin) 1% non essential amino acid 100x, 1% sodium pyruvate (Gibco), 5 10e-5 M 2-

mercaptoethanol. Before injection TC1 cells were trypsinized and washed in serum free medium.

### 1.3.3 Tumour growth:

All the animals were injected with tumor cells on day 0 and were randomized  
5 at day 7. Individual tumor growth was followed over time ( the 2 main diameters ( A, B) were measured using calipers twice a week, A x B represents the "tumor surface" and the average of the 5 values / groups is showed on a graphic over time: 6 weeks

### 1.3.4 CMI read out

#### In vitro lymphoproliferation

10 Lymphoproliferation was performed on individual spleens and on lymph node pools. 200000 spleen cells or popliteal lymph node cells were plated in triplicate, in 96 well microplate, in RPMI medium containing 1% normal mice serum and additives . After 72 hrs of in vitro re-stimulation with different amounts of PD1/3 E7 (1, 0.1, 0.01 µg/ml) or E7 ( 10-1-0.1 µg/ml ) After 72hrs, 100 µl of culture  
15 supernatant were removed and replaced by fresh medium containing 1µCi 3H thymidine (Amersham 5Ci/mmol) . After 16 hrs, cells were harvested onto filter plates. Incorporated radioactivity was counted in a β counter. Results are expressed in CPM (mean of triplicate wells) or as stimulation indexes (mean CPM in cultures with antigen / mean CPM in cultures without antigen).

20

### 1.3.5 CTL assay

20 10e6 spleen cells were co-cultured with 2 10e6 irradiated (18000r) TC1 cells (E7 expressing tumor) for 7 days in the presenced or absence of ConA sup. (2%)

25 Target cells used to assess cytotoxicity were either Cr51 ( DuPont NEN 37MBq/ml) loaded (1hr at 37°C) TC1 cells or E7 pulsed EL4 cells (for 1 hr at 37°C during the Cr 51 loading of the cells 10µg/ml of E7-derived peptide (49-57) (QCB) compared to EL4 cells NK dependant lysis was assessed on K562 target cells 2000 target cells were added / well of 96 well plate ( V botttom nunc 2-45128) with 100/1  
30 being the highest Effector / target ratio. Controls for spontaneous or maximal Cr51 release were performed in sextuplet and were targets in medium or in triton 1.5%. All plates were gently centrifuged and incubated for 4 hrs at 37 in 7% CO2. 50 µl of the

supernatant was deposited on 96w Lumaplate (Packard) let dry O/N and counted in a Top Count counter. Data is expressed as percent specific lysis which is calculated from the c.p.m. by the formula (experimental release - spontaneous release) / (maximal release - spontaneous release) X 100.

#### 5 Serology

Quantitation of anti E7 antibody was performed by Elisa using E7as coating antigen. Antigen and antibody solutions were used at 50 µl per well. Antigen was diluted at a final concentration of 3 µg/ml in carbonate buffer pH9.5 and was adsorbed overnight at 4°C to the wells of 96 wells microtiter plates (Maxisorb Immuno-plate,  
10 Nunc, Denmark). The plates were then incubated for 1hr at 37°C with PBS containing 1% bovine serum albumin and 0.1% Tween 20 (saturation buffer). Two-fold dilutions of sera (starting at 1/100 dilution) in the saturation buffer were added to the E7-coated plates and incubated for 1 hr 30 min at 37°C. The plates were washed 3 times with PBS 0.1% Tween 20 and biotin-conjugated anti-mouse IgG1, IgG2a or IgG2b or  
15 IgGtot (Amersham, UK) diluted 1/5000 in saturation buffer was added to each well and incubated for 1 hr 30 min at 37°C. After a washing step, streptavidin-biotinylated peroxidase complex (Amersham, UK) diluted 1/5000 in saturation buffer was added for an additional 30 min at 37°C. Plates were washed as above and incubated for 10 min with TMB( tetra-methyl-benzidine). The reaction was stopped with H2SO4 4N  
20 and read at 450 nm. Midpoint dilutions were calculated by SoftmaxPro (using a four parameters equation ).

#### EXAMPLE XIV

In a second experiment, the vaccine of the invention were tested to assess the significance of the backbone:

#### 25 Therapeutic experiment: protocol

- 10e6 TC1 cells , E7 expressing tumor cells : were injected subcutaneously (200µl) in the flank of immunocompetent C57BL/6 mice.
- 30 • 2 vaccinations, 7 and 14 days after the tumor challenge, with 5µg ProtD 1/3 E7 HPV16 injected intra- footpad (100 µl : 50µl / footpad) +/- CpG oligo; Oligo 1 (WD1001) as a phosphorothioate modified or the same Oligo (WD1006) but with phosphodiester linkage.

- 5 animals /group.

The tumor growth was monitored by measuring individual tumors twice a week and the mean tumor growth/ group of 5 animals is depicted in figure 5 and show the phosphorothioate modified oligonucleotides are effective in bringing about tumour regression.

#### Conclusions:

- All the animals that received 10e6 TC1 tumor cells develop a growing tumor.
- 100% of the animals vaccinated twice, 7 days apart, with the PD1/3 E7 HPV16 protein alone develop a tumor.
- 100% of the animals receiving the PD1/3 E7 protein + oligo WD1006 develop a tumor at the concentrations tested
- All the groups of animals that received the E7 protein + CpG 1001 at a concentration ranging from 10 to 200µg show tumor regression partial or complete(20-40%).

The first concentration at which this therapeutic effect on tumor regression is not fully obtained is E7+ 1µg CpG oligo 1001.

#### EXAMPLE XV

In a third series of experiments, the vaccines of the invention were evaluated in transgenic mice expressing E7 protein.

- The transgenic mouse strain has been generated by M. Parmentier and C. Ledent at the IRIBHN (ULB). (Ref: PNAS (USA) 1990, 87; 6176-6180).
- As transgenic mice live with the E7 HPV16 gene from birth, they are considered "tolerant" to this gene: E7 from HPV 16, in this situation is considered as a "self antigen".

- The expression of the transgene is driven by the thyroglobulin promoter. As Thyroglobulin is constitutively expressed only In the Thyroid, E7 is expressed in the thyroid.
- 5 • As a consequence of this expression, thyroid cells proliferate, mouse develop goiter and nodules which after 6 months to 1 year can evolve in invasive cancer.

The results (figure 6) of the experiments show that therapeutic vaccination with CpG oligonucleotide and antigen as described herein, results in a reduction of tumour  
10 growth and can induce complete tumour regression.

### Material & Methods

- 10e6 TC1 cells, E7 expressing tumor cells : were injected subcutaneously  
15 (200µl) in the flank of male or female C57BL/6 Transgenic
  - mice were vaccinated 7 and 14 days after the tumor challenge, with 5µg ProtD 1/3 E7 HPV16 injected intra- footpad (100 µl : 50µl / footpad) in the 2 presence of CpG oligonucleotide TCT CCC AGC GTG CGC CAT and two control  
20 adjuvants:,
  - 10 animals /group
- 2 and 4 weeks after the second immunization were killed and spleens or popliteal  
25 lymph.

### Conclusion

The vaccines of the invention are effective in bringing about tumour regression in  
30 HPV induced tumours.

## CLAIMS

1. A composition comprising an E6 or E7 protein or E6/E7 fusion protein from HPV optionally linked to an immunological fusion partner, and an immunomodulatory  
5 CpG oligonucleotide.
2. A composition as claimed in claim 1 wherein the fusion partner is selected from the group; protein D or a fragment thereof from *Haemophilus influenzae* B, lipoprotein D or fragment thereof from *Haemophilus influenzae* B, NS1 or fragment thereof from Influenzae Virus, and LYTA or fragment thereof from  
10 *Streptococcus Pneumoniae*.
3. A composition as claimed in claim 1 or 2 wherein the E6 or E7 proteins are derived from HPV16 or HPV18.
4. A composition as claimed in claim 1, 2 or 3 wherein the E7 protein is mutated.
5. A composition as claimed in claim 1, 2 or 3 wherein the E6 protein is mutated.
- 15 6. A composition as claimed in any of claims 1 to 5 additionally comprising a histidine tag of at least 4 histidine residues.
7. A composition as claimed herein comprising an additional HPV antigen.
8. A composition as claimed herein where the immunomodulatory CpG oligonucleotide comprises a hexamer motif: purine purine cytosine guanine pyrimidine pyrimidine.
- 20 9. A composition as claimed herein wherein the immunomodulatory CpG oligonucleotide has two or more CpG motifs.
10. A composition as claimed herein wherein the CpG oligonucleotide contains a phosphorothioate inter-nucleotide linkage.
11. A composition as claimed herein wherein the CpG oligonucleotide is selected  
25 from the group:

OLIGO 1: TCC ATG ACG TTC CTG ACG TT

OLIGO 2: TCT CCC AGC GTG CGC CAT

OLIGO 3: ACC GAT GAC GTC GCC GGT GAC GGC ACC ACG

12. A composition as claimed herein for use in medicine.
- 5 13. A method of inducing an immune response in a patient to an HPV antigen comprising administering a safe and effective amount of a composition as claimed herein.
14. A method of preventing or treating HPV induced tumours in a patient comprising administering a safe and effective amount of a composition as claimed herein.
- 10 15. A method of preparing a composition as claimed herein, comprising admixing an E6, E7 or E6/E7 fusion protein optionally linked to an immunological fusion partner, and an immunomodulatory CpG oligonucleotide.

Fig. 1

LAS97293/CGHPV97010






Fig. 2




Fig. 3




Fig. 4




Fig. 5





Fig. 6



## SEQUENCE LISTING

## (1) GENERAL INFORMATION

5

(i) APPLICANT: BRUCK, CLAUDINE

(ii) TITLE OF THE INVENTION: VACCINE

10

(iii) NUMBER OF SEQUENCES: 23

(iv) CORRESPONDENCE ADDRESS:

15

(A) ADDRESSEE: SmithKline Beecham

(B) STREET: 2 New Horizons Court, Great West Road, B

(C) CITY: Middx

(D) STATE:

(E) COUNTRY: UK

(F) ZIP: TW8 9EP

20

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette

(B) COMPUTER: IBM Compatible

(C) OPERATING SYSTEM: DOS

25

(D) SOFTWARE: FastSEQ for Windows Version 2.0

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

30

(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

35

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Dalton, Marcus J

40

(B) REGISTRATION NUMBER:

(C) REFERENCE/DOCKET NUMBER: B45124

(ix) TELECOMMUNICATION INFORMATION:

45

(A) TELEPHONE: 0181 9756348

(B) TELEFAX: 0181 9756177

(C) TELEX:

## (2) INFORMATION FOR SEQ ID NO:1:

50

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 220 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

55

(D) TOPOLOGY: linear

Protein D 1/3 E7 his

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

60 Met Asp Pro Ser Ser His Ser Ser Asn Met Ala Asn Thr Gln Met Lys  
1 5 10 15  
Ser Asp Lys Ile Ile Ala His Arg Gly Ala Ser Gly Tyr Leu Pro  
20 25 30  
Glu His Thr Leu Glu Ser Lys Ala Leu Ala Phe Ala Gln Gln Ala Asp  
35 40 45  
65 Tyr Leu Glu Gln Asp Leu Ala Met Thr Lys Asp Gly Arg Leu Val Val

```

      50      55      60
Ile His Asp His Phe Leu Asp Gly Leu Thr Asp Val Ala Lys Lys Phe
65      70      75      80
Pro His Arg His Arg Lys Asp Gly Arg Tyr Val Ile Asp Phe Thr
5      85      90      95
Leu Lys Glu Ile Gln Ser Leu Glu Met Thr Glu Asn Phe Glu Thr Met
100      105      110
Ala Met His Gly Asp Thr Pro Thr Leu His Glu Tyr Met Leu Asp Leu
115      120      125
10 Gln Pro Glu Thr Thr Asp Leu Tyr Cys Tyr Glu Gln Leu Asn Asp Ser
130      135      140
Ser Glu Glu Glu Asp Glu Ile Asp Gly Pro Ala Gly Gln Ala Glu Pro
145      150      155      160
Asp Arg Ala His Tyr Asn Ile Val Thr Phe Cys Cys Lys Cys Asp Ser
15      165      170      175
Thr Leu Arg Leu Cys Val Gln Ser Thr His Val Asp Ile Arg Thr Leu
180      185      190
Glu Asp Leu Leu Met Gly Thr Leu Gly Ile Val Cys Pro Ile Cys Ser
195      200      205
20 Gln Lys Pro Thr Ser Gly His His His His His His
210      215      220

```

## (2) INFORMATION FOR SEQ ID NO:2:

## 25 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 663 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 Protein D 1/3 E7 his

30

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

ATGGATCCAA GCAGCCATTC ATCAATATG GCGAATACCC AAATGAAATC AGACAAAATC
35 60
ATTATTGCTC ACCGTGGTGC TAGCGGTTAT TTACCAGAGC ATACGTTAGA ATCTAAAGCA
120
CTTGCGTTTG CACAACAGGC TGATTATTTA GAGCAAGATT TAGCAATGAC TAAGGATGGT
180
CGTTTAGTGG TTATTCACGA TCACTTTTTA GATGGCTTGA CTGATGTTGC GAAAAAATTC
40 240
CCACATCGTC ATCGTAAAGA TGGCCGTAC TATGTCATCG ACTTTACCTT AAAAGAAAT
300
CAAAGTTTAG AAATGACAGA AAACTTTGAA ACCATGGCCA TGCATGGAGA TACACCTACA
45 360
TTGCATGAAT ATATGTTAGA TTTGCAACCA GAGACAACCT ATCTCTACTG TTATGAGCAA
420
TTAAATGACA GCTCAGAGGA GGAGGATGAA ATAGATGGTC CAGCTGGACA AGCAGAACCG
480
GACAGAGCCC ATTACAATAT TGTAACTTTT TGTGCAAGT GTGACTCTAC GCTTCGGTTG
50 540
TGCGTACAAA GCACACACGT AGACATTCGT ACTTTGGAAG ACCTGTTAAT GGGCACACTA
600
GGAATTGTGT GCCCATCTG TTCTCAGAAA CCAACTAGTG GCCACCATCA CCATCACCAT
55 660
TAA
663

```

## (2) INFORMATION FOR SEQ ID NO:3:

60

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 822 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 Protein D 1/3 E6 His/HPV 16

65

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

5 60 ATGGATCCAA GCAGCCATTC ATCAAATATG GCGAATACCC AAATGAAATC AGACAAAATC  
 120 ATTATTGCTC ACCGTGGTGC TAGCGGTTAT TTACCAGAGC ATACGTTAGA ATCTAAAGCA  
 180 CTTGCGTTTG CACAACAGGC TGATTATTTA GAGCAAGATT TAGCAATGAC TAAGGATGGT  
 10 240 CGTTTAGTGG TTATTCACGA TCACTTTTGA GATGGCTTGA CTGATGTTGC GAAAAAATTC  
 300 CCACATCGTC ATCGTAAAGA TGGCCGTTAC TATGTCATCG ACTTTACCTT AAAAGAAATT  
 360 CAAAGTTTAG AAATGACAGA AAAGTTTGA ACCATGGCCA TGTTTCAGGA CCCACAGGAG  
 15 420 CGACCCAGAA AGTTACCACA GTTATGCACA GAGCTGCAAA CAACTATACA TGATATAATA  
 480 TTAGAATGTG TGTACTGCAA GCAACAGTTA CTGCGACGTG AGGTATATGA CTTTGCTTTT  
 20 540 CGGGATTTAT GCATAGTATA TAGAGATGGG AATCCATATG CTGTATGTGA TAAATGTTTA  
 600 AAGTTTTATT CTAATAATTAG TGAGTATAGA CATTATTGTT ATAGTTTGTA TGGAAACAACA  
 660 TTAGAACAGC AATACAACAA ACCGTTGTGT GATTGTGTTA TTAGGTGTAT TAACTGTCAA  
 25 720 AAGCCACTGT GTCCTGAAGA AAAGCAAAGA CATCTGGACA AAAAGCAAAG ATTCCATAAT  
 780 ATAAGGGGTC GGTGGACCGG TCGATGTATG TCTTGTTGCA GATCATCAAG AACACGTAGA  
 30 822 GAAACCCAGC TGACTAGTGG CCACCATCAC CATCACCATT AA

## (2) INFORMATION FOR SEQ ID NO:4:

35 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 274 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 40 Protein D 1/3 E6 His/HPV 16

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

45 Met Asp Pro Ser Ser His Ser Ser Asn Met Ala Asn Thr Gln Met Lys  
 1 5 10 15  
 Ser Asp Lys Ile Ile Ile Ala His Arg Gly Ala Ser Gly Tyr Leu Pro  
 20 25 30  
 Glu His Thr Leu Glu Ser Lys Ala Leu Ala Phe Ala Gln Gln Ala Asp  
 35 40 45  
 50 Tyr Leu Glu Gln Asp Leu Ala Met Thr Lys Asp Gly Arg Leu Val Val  
 50 55 60  
 Ile His Asp His Phe Leu Asp Gly Leu Thr Asp Val Ala Lys Lys Phe  
 65 70 75 80  
 Pro His Arg His Arg Lys Asp Gly Arg Tyr Tyr Val Ile Asp Phe Thr  
 85 90 95  
 55 Leu Lys Glu Ile Gln Ser Leu Glu Met Thr Glu Asn Phe Glu Thr Met  
 100 105 110  
 Ala Met Phe Gln Asp Pro Gln Glu Arg Pro Arg Lys Leu Pro Gln Leu  
 115 120 125  
 60 Cys Thr Glu Leu Gln Thr Thr Ile His Asp Ile Ile Leu Glu Cys Val  
 130 135 140  
 Tyr Cys Lys Gln Gln Leu Leu Arg Arg Glu Val Tyr Asp Phe Ala Phe  
 145 150 155 160  
 Arg Asp Leu Cys Ile Val Tyr Arg Asp Gly Asn Pro Tyr Ala Val Cys  
 165 170 175  
 65 Asp Lys Cys Leu Lys Phe Tyr Ser Lys Ile Ser Glu Tyr Arg His Tyr



				180						185					190	
	Cys	Tyr	Ser	Leu	Tyr	Gly	Thr	Thr	Leu	Glu	Gln	Gln	Tyr	Asn	Lys	Pro
			195					200					205			
	Leu	Cys	Asp	Leu	Leu	Ile	Arg	Cys	Ile	Asn	Cys	Gln	Lys	Pro	Leu	Cys
5		210					215					220				
	Pro	Glu	Glu	Lys	Gln	Arg	His	Leu	Asp	Lys	Lys	Gln	Arg	Phe	His	Asn
	225					230					235					240
	Ile	Arg	Gly	Arg	Trp	Thr	Gly	Arg	Cys	Met	Ser	Cys	Cys	Arg	Ser	Ser
					245					250					255	
10	Arg	Thr	Arg	Arg	Glu	Thr	Gln	Leu	Thr	Ser	Gly	His	His	His	His	His
				260					265					270		
	His															

15 (2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1116 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

Protein: D 1/3 E6/E7/ HPV16

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

25 ATGGATCCAA GCAGCCATTC ATCAAATATG GCGAATACCC AAATGAAATC AGACAAAATC  
60  
ATTATTGCTC ACCGTGGTGC TAGCGGTTAT TTACCAGAGC ATACGTTAGA ATCTAAAGCA  
120  
30 CTTGCGTTTG CACAACAGGC TGATTATTTA GAGCAAGATT TAGCAATGAC TAAGGATGGT  
180  
CGTTTAGTGG TTATTCACGA TCACTTTTTA GATGGCTTGA CTGATGTTGC GAAAAAATTC  
240  
CCACATCGTC ATCGTAAAGA TGGCCGTTAC TATGTCATCG ACTTTACCTT AAAAGAAATC  
35 300  
CAAAGTTTAG AAATGACAGA AACTTTGAA ACCATGGCCA TGTTTCAGGA CCCACAGGAG  
360  
CGACCCAGAA AGTTACCACA GTTATGCACA GAGCTGCAAA CAACTATACA TGATATAATA  
420  
40 TTAGAATGTG TGTACTGCAA GCAACAGTTA CTGCGACGTG AGGTATATGA CTTTGCTTTT  
480  
CGGGATTAT GCATAGTATA TAGAGATGGG AATCCATATG CTGTATGTGA TAAATGTTTA  
540  
AAGTTTTATT CTAATAATTAG TGAGTATAGA CATTATTGTT ATAGTTTGTA TGAACAACA  
45 600  
TTAGAACAGC AATACAACAA ACCGTTGTGT GATTTGTAA TTAGGTGTAT TAACTGTCAA  
660  
AAGCCACTGT GTCCTGAAGA AAAGCAAAGA CATCTGGACA AAAAGCAAAG ATTCCATAAT  
720  
50 ATAAGGGGTC GGTGGACCGG TCGATGTATG TCTTGTGCA GATCATCAAG AACACGTAGA  
780  
GAAACCCAGC TGATGCATGG AGATACACCT ACATTGCATG AATATATGTT AGATTGCAA  
840  
CCAGAGACAA CTGATCTCTA CTGTTATGAG CAATTAAATG ACAGCTCAGA GGAGGAGGAT  
55 900  
GAAATAGATG GTCCAGCTGG ACAAGCAGAA CCGGACAGAG CCCATTACAA TATTGTAACC  
960  
TTTTGTTGCA AGTGTGACTC TACGCTTCGG TTGTGCGTAC AAAGCACACA CGTAGACATT  
1020  
60 CGTACTTTGG AAGACCTGTT AATGGGCACA CTAGGAATTG TGTGCCCCAT CTGTTCTCAG  
1080  
AAACCAACTA GTGGCCACCA TCACCATCAC CATTAA  
1116

65 (2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 372 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 Protein D 1/3 E6/E7/ HPV16

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

10 Met Asp Pro Ser Ser His Ser Ser Asn Met Ala Asn Thr Gln Met Lys
    1      5      10      15
    Ser Asp Lys Ile Ile Ala His Arg Gly Ala Ser Gly Tyr Leu Pro
        20      25      30
    Glu His Thr Leu Glu Ser Lys Ala Leu Ala Phe Ala Gln Ala Asp
        35      40      45
15 Tyr Leu Glu Gln Asp Leu Ala Met Thr Lys Asp Gly Arg Leu Val Val
    50      55      60
    Ile His Asp His Phe Leu Asp Gly Leu Thr Asp Val Ala Lys Lys Phe
    65      70      75      80
20 Pro His Arg His Arg Lys Asp Gly Arg Tyr Tyr Val Ile Asp Phe Thr
    85      90      95
    Leu Lys Glu Ile Gln Ser Leu Glu Met Thr Glu Asn Phe Glu Thr Met
    100      105      110
    Ala Met Phe Gln Asp Pro Gln Glu Arg Pro Arg Lys Leu Pro Gln Leu
    115      120      125
25 Cys Thr Glu Leu Gln Thr Thr Ile His Asp Ile Ile Leu Glu Cys Val
    130      135      140
    Tyr Cys Lys Gln Gln Leu Leu Arg Arg Glu Val Tyr Asp Phe Ala Phe
    145      150      155      160
30 Arg Asp Leu Cys Ile Val Tyr Arg Asp Gly Asn Pro Tyr Ala Val Cys
    165      170      175
    Asp Lys Cys Leu Lys Phe Tyr Ser Lys Ile Ser Glu Tyr Arg His Tyr
    180      185      190
    Cys Tyr Ser Leu Tyr Gly Thr Thr Leu Glu Gln Gln Tyr Asn Lys Pro
    195      200      205
35 Leu Cys Asp Leu Leu Ile Arg Cys Ile Asn Cys Gln Lys Pro Leu Cys
    210      215      220
    Pro Glu Glu Lys Gln Arg His Leu Asp Lys Lys Gln Arg Phe His Asn
    225      230      235      240
40 Ile Arg Gly Arg Trp Thr Gly Arg Cys Met Ser Cys Cys Arg Ser Ser
    245      250      255
    Arg Thr Arg Arg Glu Thr Gln Leu Met His Gly Asp Thr Pro Thr Leu
    260      265      270
    His Glu Tyr Met Leu Asp Leu Gln Pro Glu Thr Thr Asp Leu Tyr Cys
    275      280      285
45 Tyr Glu Gln Leu Asn Asp Ser Ser Glu Glu Glu Asp Glu Ile Asp Gly
    290      295      300
    Pro Ala Gly Gln Ala Glu Pro Asp Arg Ala His Tyr Asn Ile Val Thr
    305      310      315      320
50 Phe Cys Cys Lys Cys Asp Ser Thr Leu Arg Leu Cys Val Gln Ser Thr
    325      330      335
    His Val Asp Ile Arg Thr Leu Glu Asp Leu Leu Met Gly Thr Leu Gly
    340      345      350
    Ile Val Cys Pro Ile Cys Ser Gln Lys Pro Thr Ser Gly His His His
    355      360      365
55 His His His
    370
  
```

(2) INFORMATION FOR SEQ ID NO:7:

60

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 663 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 Protein D 1/3 E7 mutated HPV 16

65

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

5 60 ATGGATCCAA GCAGCCATTC ATCAAATATG GCGAATACCC AAATGAAATC AGACAAAATC  
 ATTATTGCTC ACCGTGGTGC TAGCGGTAT TTACCAGAGC ATACGTTAGA ATCTAAAGCA  
 120 CTTGCGTTTG CACAACAGGC TGATTATTTA GAGCAAGATT TAGCAATGAC TAAGGATGGT  
 180 CGTTTAGTGG TTATTCACGA TCACTTTTTA GATGGCTTGA CTGATGTTGC GAAAAAATTC  
 240 CCACATCGTC ATCGTAAAGA TGGCCGTTAC TATGTCATCG ACTTTACCTT AAAAGAAATT  
 300 CAAAGTTTAG AAATGACAGA AAACCTTTGAA ACCATGGCCA TGCATGGAGA TACACCTACA  
 360 TTGCATGAAT ATATGTTAGA TTTGCAACCA GAGACAACCTG ATCTCTACGG TTATCAGCAA  
 420 TTAAATGACA GCTCAGAGGA GGAGGATGAA ATAGATGGTC CAGCTGGACA AGCAGAACCG  
 480 GACAGAGCCC ATTACAATAT TGTAACCTTT TGTGCAAGT GTGACTCTAC GCTTCGGTTG  
 20 540 TGGCTACAAA GCACACACGT AGACATTCGT ACTTTGGAAG ACCTGTTAAT GGGCACACTA  
 600 GGAATTGTGT GCCCCATCTG TTCTCAGAAA CCAACTAGTG GCCACCATCA CCATCACCAT  
 25 660 TAA  
 663

## (2) INFORMATION FOR SEQ ID NO:8:

- 30 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 220 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 35 (D) TOPOLOGY: linear

Protein D 1/3 E7 mutated HPV 16

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

40 Met Asp Pro Ser Ser His Ser Ser Asn Met Ala Asn Thr Gln Met Lys  
 1 5 10 15  
 Ser Asp Lys Ile Ile Ile Ala His Arg Gly Ala Ser Gly Tyr Leu Pro  
 20 25 30  
 45 Glu His Thr Leu Glu Ser Lys Ala Leu Ala Phe Ala Gln Ala Asp  
 35 40 45  
 Tyr Leu Glu Gln Asp Leu Ala Met Thr Lys Asp Gly Arg Leu Val Val  
 50 55 60  
 Ile His Asp His Phe Leu Asp Gly Leu Thr Asp Val Ala Lys Lys Phe  
 65 70 75 80  
 50 Pro His Arg His Arg Lys Asp Gly Arg Tyr Tyr Val Ile Asp Phe Thr  
 85 90 95  
 Leu Lys Glu Ile Gln Ser Leu Glu Met Thr Glu Asn Phe Glu Thr Met  
 100 105 110  
 Ala Met His Gly Asp Thr Pro Thr Leu His Glu Tyr Met Leu Asp Leu  
 115 120 125  
 55 Gln Pro Glu Thr Thr Asp Leu Tyr Gly Tyr Gln Gln Leu Asn Asp Ser  
 130 135 140  
 Ser Glu Glu Glu Asp Glu Ile Asp Gly Pro Ala Gly Gln Ala Glu Pro  
 145 150 155 160  
 60 Asp Arg Ala His Tyr Asn Ile Val Thr Phe Cys Cys Lys Cys Asp Ser  
 165 170 175  
 Thr Leu Arg Leu Cys Val Gln Ser Thr His Val Asp Ile Arg Thr Leu  
 180 185 190  
 Glu Asp Leu Leu Met Gly Thr Leu Gly Ile Val Cys Pro Ile Cys Ser  
 195 200 205  
 65 Gln Lys Pro Thr Ser Gly His His His His His His

210

215

220

## (2) INFORMATION FOR SEQ ID NO:9:

- 5 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 879 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 10 CLYTA E6 His HPV 16

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

15 ATGAAAGGGG GAATTGTACA TTCAGACGGC TCTTATCCAA AAGACAAGTT TGAGAAAATC  
 60 AATGGCACTT GGTACTACTT TGACAGTTCA GGCTATATGC TTGCAGACCG CTGGAGGAAG  
 120 CACACAGACG GCAACTGGTA CTGGTTCGAC AACTCAGGCG AAATGGCTAC AGGCTGGAAG  
 180 AAAATCGCTG ATAAGTGGTA CTATTCAAC GAAGAAGGTG CCATGAAGAC AGGCTGGGTC  
 20 240 AAGTACAAGG ACACTTGGTA CTAATTAGAC GCTAAAGAAG GCGCCATGGT ATCAAATGCC  
 300 TTTATCCAGT CAGCGGACGG AACAGGCTGG TACTACCTCA AACCAGACGG AACACTGGCA  
 25 360 GACAGGCCAG AATTGGCCAG CATGCTGGAC ATGGCCATGT TTCAGGACCC ACAGGAGCGA  
 420 CCCAGAAAGT TACCACAGTT ATGCACAGAG CTGCAAACAA CTATACATGA TATAATATTA  
 480 GAATGTGTGT ACTGCAAGCA ACAGTTACTG CGACGTGAGG TATATGACTT TGCTTTTCGG  
 30 540 GATTTATGCA TAGTATATAG AGATGGGAAT CCATATGCTG TATGTGATAA ATGTTTAAAG  
 600 TTTTATTCTA AAATTAGTGA GTATAGACAT TATTGTTATA GTTTGTATGG AACACATTA  
 35 660 GAACAGCAAT ACAACAAACC GTTGTGTGAT TTGTTAATTA GGTGTATTAA CTGTCAAAG  
 720 CCACTGTGTC CTGAAGAAAA GCAAAGACAT CTGGACAAAA AGCAAAGATT CCATAATATA  
 780 AGGGGTCGGT GGACCGGTCG ATGTATGTCT TGTTCAGAT CATCAAGAAC ACGTAGAGAA  
 40 840 ACCCAGCTGA CTAGTGGCCA CCATCACCAT CACCATTAA  
 879

## (2) INFORMATION FOR SEQ ID NO:10:

- 45 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 293 amino acids  
 (B) TYPE: amino acid  
 50 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 CLYTA E6 His HPV 16

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

55 Met Lys Gly Gly Ile Val His Ser Asp Gly Ser Tyr Pro Lys Asp Lys  
 1 5 10 15  
 Phe Glu Lys Ile Asn Gly Thr Trp Tyr Tyr Phe Asp Ser Ser Gly Tyr  
 20 25 30  
 60 Met Leu Ala Asp Arg Trp Arg Lys His Thr Asp Gly Asn Trp Tyr Trp  
 35 40 45  
 Phe Asp Asn Ser Gly Glu Met Ala Thr Gly Trp Lys Lys Ile Ala Asp  
 50 55 60  
 Lys Trp Tyr Tyr Phe Asn Glu Glu Gly Ala Met Lys Thr Gly Trp Val  
 65 65 70 75 80  
 Lys Tyr Lys Asp Thr Trp Tyr Tyr Leu Asp Ala Lys Glu Gly Ala Met

85 90 95  
 Val Ser Asn Ala Phe Ile Gln Ser Ala Asp Gly Thr Gly Trp Tyr Tyr  
 100 105 110  
 5 Leu Lys Pro Asp Gly Thr Leu Ala Asp Arg Pro Glu Leu Ala Ser Met  
 115 120 125  
 Leu Asp Met Ala Met Phe Gln Asp Pro Gln Glu Arg Pro Arg Lys Leu  
 130 135 140  
 Pro Gln Leu Cys Thr Glu Leu Gln Thr Thr Ile His Asp Ile Ile Leu  
 145 150 155 160  
 10 Glu Cys Val Tyr Cys Lys Gln Gln Leu Leu Arg Arg Glu Val Tyr Asp  
 165 170 175  
 Phe Ala Phe Arg Asp Leu Cys Ile Val Tyr Arg Asp Gly Asn Pro Tyr  
 180 185 190  
 Ala Val Cys Asp Lys Cys Leu Lys Phe Tyr Ser Lys Ile Ser Glu Tyr  
 195 200 205  
 15 Arg His Tyr Cys Tyr Ser Leu Tyr Gly Thr Thr Leu Glu Gln Gln Tyr  
 210 215 220  
 Asn Lys Pro Leu Cys Asp Leu Leu Ile Arg Cys Ile Asn Cys Gln Lys  
 225 230 235 240  
 20 Pro Leu Cys Pro Glu Lys Gln Arg His Leu Asp Lys Lys Gln Arg  
 245 250 255  
 Phe His Asn Ile Arg Gly Arg Trp Thr Gly Arg Cys Met Ser Cys Cys  
 260 265 270  
 Arg Ser Ser Arg Thr Arg Arg Glu Thr Gln Leu Thr Ser Gly His His  
 275 280 285  
 25 His His His His  
 290

## (2) INFORMATION FOR SEQ ID NO:11:

30

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 720 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

35

CLYTA E7 HIS HPV 16

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

40 ATGAAAGGGG GAATTGTACA TTCAGACGGC TCTTATCCAA AAGACAAGTT TGAGAAAATC  
 60 AATGGCACTT GGTACTACTT TGACAGTTCA GGCTATATGC TTGCAGACCG CTGGAGGAAG  
 120 CACACAGACG GCAACTGGTA CTGGTTCGAC AACTCAGGCG AAATGGCTAC AGGCTGGAAG  
 45 180 AAAATCGCTG ATAAGTGGTA CTATTTC AAC GAAGAAGGTG CCATGAAGAC AGGCTGGGTC  
 240 AAGTACAAGG AACTTGGTA CTA CTACTTAGAC GCTAAAGAAG GCGCCATGGT ATCAAATGCC  
 300 TTTATCCAGT CAGCGGACGG AACAGGCTGG TACTACCTCA AACCAGACGG AACACTGGCA  
 50 360 GACAGGCCAG AATTGGCCAG CATGCTGGAC ATGGCCATGC ATGGAGATAC ACCTACATTG  
 420 CATGAATATA TGTTAGATTT GCAACCAGAG ACAACTGATC TCTACTGTTA TGAGCAATTA  
 55 480 AATGACAGCT CAGAGGAGGA GGATGAAATA GATGGTCCAG CTGGACAAGC AGAACCGGAC  
 540 AGAGCCCATT ACAATATTGT AACCTTTTGT TGCAAGTGTG ACTCTACGCT TCGGTTGTGC  
 600 GTACAAAGCA CACACGTAGA CATTCTGACT TTGGAAGACC TGTTAATGGG CACACTAGGA  
 660 ATTGTGTGCC CCATCTGTTC TCAGAAACCA ACTAGTGGCC ACCATCACCA TCACCATTAA  
 720

65

## (2) INFORMATION FOR SEQ ID NO:12:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 240 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

CLYTA E7 HIS HPV 16

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

```

10 Met Lys Gly Gly Ile Val His Ser Asp Gly Ser Tyr Pro Lys Asp Lys
    1      5      10      15
    Phe Glu Lys Ile Asn Gly Thr Trp Tyr Tyr Phe Asp Ser Ser Gly Tyr
    20      25      30
    Met Leu Ala Asp Arg Trp Arg Lys His Thr Asp Gly Asn Trp Tyr Trp
    35      40      45
15 Phe Asp Asn Ser Gly Glu Met Ala Thr Gly Trp Lys Lys Ile Ala Asp
    50      55      60
    Lys Trp Tyr Tyr Phe Asn Glu Glu Gly Ala Met Lys Thr Gly Trp Val
    65      70      75      80
20 Lys Tyr Lys Asp Thr Trp Tyr Tyr Leu Asp Ala Lys Glu Gly Ala Met
    85      90      95
    Val Ser Asn Ala Phe Ile Gln Ser Ala Asp Gly Thr Gly Trp Tyr Tyr
    100      105      110
    Leu Lys Pro Asp Gly Thr Leu Ala Asp Arg Pro Glu Leu Ala Ser Met
    115      120      125
25 Leu Asp Met Ala Met His Gly Asp Thr Pro Thr Leu His Glu Tyr Met
    130      135      140
    Leu Asp Leu Gln Pro Glu Thr Thr Asp Leu Tyr Cys Tyr Glu Gln Leu
    145      150      155      160
30 Asn Asp Ser Ser Glu Glu Asp Glu Ile Asp Gly Pro Ala Gly Gln
    165      170      175
    Ala Glu Pro Asp Arg Ala His Tyr Asn Ile Val Thr Phe Cys Cys Lys
    180      185      190
    Cys Asp Ser Thr Leu Arg Leu Cys Val Gln Ser Thr His Val Asp Ile
    195      200      205
35 Arg Thr Leu Glu Asp Leu Leu Met Gly Thr Leu Gly Ile Val Cys Pro
    210      215      220
    Ile Cys Ser Gln Lys Pro Thr Ser Gly His His His His His His
    225      230      235

```

## (2) INFORMATION FOR SEQ ID NO:13:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1173 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

CLYTA E6E7 His HPV16

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

```

    ATGAAAGGGG GAATTGTACA TTCAGACGGC TCTTATCCAA AAGACAAGTT TGAGAAAATC
60
    AATGGCACTT GGTACTACTT TGACAGTTCA GGCTATATGC TTGCAGACCG CTGGAGGAAG
55 120
    CACACAGACG GCAACTGGTA CTGGTTCGAC AACTCAGGCG AAATGGCTAC AGGCTGGAAG
    180
    AAAATCGCTG ATAAGTGGTA CTATTTCAAC GAAGAAGGTG CCATGAAGAC AGGCTGGGTC
    240
60 AAGTACAAGG ACACTTGGTA CTA CTAGTAC GCTAAAGAAG GCGCCATGGT ATCAAATGCC
    300
    TTTATCCAGT CAGCGGACGG AACAGGCTGG TACTACCTCA AACCAGACGG AACACTGGCA
    360
65 GACAGGCCAG AATTGGCCAG CATGCTGGAC ATGGCCATGT TTCAGGACCC ACAGGAGCGA
    420

```

480 CCCAGAAAGT TACCACAGTT ATGCACAGAG CTGCAAACAA CTATACATGA TATAATATTA  
 540 GAATGTGTGT ACTGCAAGCA ACAGTTACTG CGACGTGAGG TATATGACTT TGCTTTTCGG  
 5 540 GATTTATGCA TAGTATATAG AGATGGGAAT CCATATGCTG TATGTGATAA ATGTTTAAAG  
 600 TTTTATTCTA AAATTAGTGA GTATAGACAT TATTGTTATA GTTTGTATGG AACAAACATTA  
 660 GAACAGCAAT ACAACAAACC GTTGTGTGAT TTGTTAATTA GGTGTATTAA CTGTCAAAAG  
 10 720 CCACTGTGTC CTGAAGAAAA GCAAAGACAT CTGGACAAAA AGCAAAGATT CCATAATATA  
 780 AGGGGTCGGT GGACCGGTCG ATGTATGTCT TGTTGCAGAT CATCAAGAAC ACGTAGAGAA  
 840 ACCCAGCTGA TGCATGGAGA TACACCTACA TTGCATGAAT ATATGTTAGA TTTGCAACCA  
 15 900 GAGACAACCTG ATCTCTACTG TTATGAGCAA TTAAATGACA GCTCAGAGGA GGAGGATGAA  
 960 ATAGATGGTC CAGCTGGACA AGCAGAACCG GACAGAGCCC ATTACAATAT TGTAACCTTT  
 20 1020 TGTTCGAAGT GTGACTCTAC GCTTCGGTTG TCGGTACAAA GCACACACGT AGACATTCGT  
 1080 ACTTTGGAAG ACCTGTTAAT GGGCACACTA GGAATTGTGT GCCCATCTG TTCTCAGAAA  
 1140 CCAACTAGTG GCCACCATCA CCATCACCAT TAA  
 25 1173

## (2) INFORMATION FOR SEQ ID NO:14:

- 30 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 391 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 35 CLYTA E6E7 His HPV16

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

40 Met Lys Gly Gly Ile Val His Ser Asp Gly Ser Tyr Pro Lys Asp Lys  
 1 5 10 15  
 Phe Glu Lys Ile Asn Gly Thr Trp Tyr Tyr Phe Asp Ser Ser Gly Tyr  
 20 25 30  
 Met Leu Ala Asp Arg Trp Arg Lys His Thr Asp Gly Asn Trp Tyr Trp  
 35 40 45  
 45 Phe Asp Asn Ser Gly Glu Met Ala Thr Gly Trp Lys Lys Ile Ala Asp  
 50 55 60  
 Lys Trp Tyr Tyr Phe Asn Glu Glu Gly Ala Met Lys Thr Gly Trp Val  
 65 70 75 80  
 Lys Tyr Lys Asp Thr Trp Tyr Tyr Leu Asp Ala Lys Glu Gly Ala Met  
 85 90 95  
 50 Val Ser Asn Ala Phe Ile Gln Ser Ala Asp Gly Thr Gly Trp Tyr Tyr  
 100 105 110  
 Leu Lys Pro Asp Gly Thr Leu Ala Asp Arg Pro Glu Leu Ala Ser Met  
 115 120 125  
 55 Leu Asp Met Ala Met Phe Gln Asp Pro Gln Glu Arg Pro Arg Lys Leu  
 130 135 140  
 Pro Gln Leu Cys Thr Glu Leu Gln Thr Thr Ile His Asp Ile Ile Leu  
 145 150 155 160  
 Glu Cys Val Tyr Cys Lys Gln Gln Leu Leu Arg Arg Glu Val Tyr Asp  
 165 170 175  
 60 Phe Ala Phe Arg Asp Leu Cys Ile Val Tyr Arg Asp Gly Asn Pro Tyr  
 180 185 190  
 Ala Val Cys Asp Lys Cys Leu Lys Phe Tyr Ser Lys Ile Ser Glu Tyr  
 195 200 205  
 65 Arg His Tyr Cys Tyr Ser Leu Tyr Gly Thr Thr Leu Glu Gln Gln Tyr  
 210 215 220

Asn Lys Pro Leu Cys Asp Leu Leu Ile Arg Cys Ile Asn Cys Gln Lys  
 225 230 235 240  
 Pro Leu Cys Pro Glu Glu Lys Gln Arg His Leu Asp Lys Lys Gln Arg  
 245 250 255  
 5 Phe His Asn Ile Arg Gly Arg Trp Thr Gly Arg Cys Met Ser Cys Cys  
 260 265 270  
 Arg Ser Ser Arg Thr Arg Arg Glu Thr Gln Leu Met His Gly Asp Thr  
 275 280 285  
 Pro Thr Leu His Glu Tyr Met Leu Asp Leu Gln Pro Glu Thr Thr Asp  
 290 295 300  
 10 Leu Tyr Cys Tyr Glu Gln Leu Asn Asp Ser Ser Glu Glu Glu Asp Glu  
 305 310 315 320  
 Ile Asp Gly Pro Ala Gly Gln Ala Glu Pro Asp Arg Ala His Tyr Asn  
 325 330 335  
 15 Ile Val Thr Phe Cys Cys Lys Cys Asp Ser Thr Leu Arg Leu Cys Val  
 340 345 350  
 Gln Ser Thr His Val Asp Ile Arg Thr Leu Glu Asp Leu Leu Met Gly  
 355 360 365  
 Thr Leu Gly Ile Val Cys Pro Ile Cys Ser Gln Lys Pro Thr Ser Gly  
 370 375 380  
 20 His His His His His His  
 385 390

## (2) INFORMATION FOR SEQ ID NO:15:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 684 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

Protein D 1/3 E7 his HPV 18

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

35 ATGGATCCAA GCAGCCATTC ATCAAATATG GCGAATACCC AAATGAAATC AGACAAAATC  
 60 ATTATTGCTC ACCGTGGTGC TAGCGGTTAT TTACCAGAGC ATACGTTAGA ATCTAAAGCA  
 120 CTTGCGTTTG CACAACAGGC TGATTATTTA GAGCAAGATT TAGCAATGAC TAAGGATGGT  
 40 180 CGTTTAGTGG TTATTCACGA TCACTTTTGA GATGGCTTGA CTGATGTTGC GAAAAAATTC  
 240 CCACATCGTC ATCGTAAAGA TGGCCGTTAC TATGTCATCG ACTTTACCTT AAAAGAAATT  
 45 300 CAAAGTTTAG AAATGACAGA AAACCTTGAA ACCATGGCCA TGCATGGACC TAAGGCAACA  
 360 TTGCAAGACA TTGTATTGCA TTTAGAGCCC CAAAATGAAA TTCCGGTTGA CCTTCTATGT  
 420 CACGAGCAAT TAAGCGACTC AGAGGAAGAA AACGATGAAA TAGATGAAGT TAATCATCAA  
 50 480 CATTACCAG CCCGACGAGC CGAACCACAA CGTCACACAA TGTTGTGTAT GTGTTGTAAG  
 540 TGTGAAGCCA GAATTGAGCT AGTAGTAGAA AGCTCAGCAG ACGACCTTCG AGCATTCCAG  
 55 600 CAGCTGTTTC TGAACACCCT GTCCTTTGTG TGTCGGTGGT GTGCATCCCA GCAGACTAGT  
 660 GGCCACCATC ACCATCACCA TTAA  
 684

## (2) INFORMATION FOR SEQ ID NO:16:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 228 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single



(D) TOPOLOGY: linear  
Protein D 1/3 E7 his HPV 18

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

```

5  Met Asp Pro Ser Ser His Ser Ser Asn Met Ala Asn Thr Gln Met Lys
   1      5      10      15
   Ser Asp Lys Ile Ile Ile Ala His Arg Gly Ala Ser Gly Tyr Leu Pro
      20      25      30
10  Glu His Thr Leu Glu Ser Lys Ala Leu Ala Phe Ala Gln Gln Ala Asp
      35      40      45
   Tyr Leu Glu Gln Asp Leu Ala Met Thr Lys Asp Gly Arg Leu Val Val
      50      55      60
   Ile His Asp His Phe Leu Asp Gly Leu Thr Asp Val Ala Lys Lys Phe
      65      70      75      80
15  Pro His Arg His Arg Lys Asp Gly Arg Tyr Tyr Val Ile Asp Phe Thr
      85      90      95
   Leu Lys Glu Ile Gln Ser Leu Glu Met Thr Glu Asn Phe Glu Thr Met
      100      105      110
20  Ala Met His Gly Pro Lys Ala Thr Leu Gln Asp Ile Val Leu His Leu
      115      120      125
   Glu Pro Gln Asn Glu Ile Pro Val Asp Leu Leu Cys His Glu Gln Leu
      130      135      140
   Ser Asp Ser Glu Glu Glu Asn Asp Glu Ile Asp Glu Val Asn His Gln
      145      150      155      160
25  His Leu Pro Ala Arg Arg Ala Glu Pro Gln Arg His Thr Met Leu Cys
      165      170      175
   Met Cys Cys Lys Cys Glu Ala Arg Ile Glu Leu Val Val Glu Ser Ser
      180      185      190
30  Ala Asp Asp Leu Arg Ala Phe Gln Gln Leu Phe Leu Asn Thr Leu Ser
      195      200      205
   Phe Val Cys Pro Trp Cys Ala Ser Gln Gln Thr Ser Gly His His His
      210      215      220
   His His His
      225
35

```

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 110 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
Thioredoxin

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

```

Met Ser Asp Lys Ile Ile His Leu Thr Asp Asp Ser Phe Asp Thr Asp
1      5      10      15
50 Val Leu Lys Ala Asp Gly Ala Ile Leu Val Asp Phe Trp Ala Glu Trp
      20      25      30
   Cys Gly Pro Cys Lys Met Ile Ala Pro Ile Leu Asp Glu Ile Ala Asp
      35      40      45
   Glu Tyr Gln Gly Lys Leu Thr Val Ala Lys Leu Asn Ile Asp Gln Asn
      50      55      60
55 Pro Gly Thr Ala Pro Lys Tyr Gly Ile Arg Gly Ile Pro Thr Leu Leu
      65      70      75      80
   Leu Phe Lys Asn Gly Glu Val Ala Ala Thr Lys Val Gly Ala Leu Ser
      85      90      95
60 Lys Gly Gln Leu Lys Glu Phe Leu Asp Ala Asn Leu Ala
      100      105

```

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 684 base pairs

(B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 Protein D 1/3 E7 mutated HPV 18

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

```

ATGGATCCAA GCAGCCATTC ATCAATATG GCGAATACCC AAATGAAATC AGACAAAATC
60
ATTATTGCTC ACCGTGGTGC TAGCGGTTAT TTACCAGAGC ATACGTTAGA ATCTAAAGCA
10 120
CTTGCGTTTG CACAACAGGC TGATTATTTA GAGCAAGATT TAGCAATGAC TAAGGATGGT
180
CGTTTAGTGG TTATTCACGA TCACTTTTGA GATGGCTTGA CTGATGTTGC GAAAAAATTC
15 240
CCACATCGTC ATCGTAAAGA TGGCCGTAC TATGTCATCG ACTTTACCTT AAAAGAAATT
300
CAAAGTTTAG AAATGACAGA AAACCTTGAA ACCATGGCCA TGCATGGACC TAAGGCAACA
360
TTGCAAGACA TTGTATTGCA TTTAGAGCCC CAAAATGAAA TTCCGGTTGA CCTTCTAGGT
20 420
CACCAGCAAT TAAGCGACTC AGAGGAAGAA AACGATGAAA TAGATGGAGT TAATCATCAA
480
CATTTACCAG CCCGACGAGC CGAACCACAA CGTCACACAA TGTTGTGTAT GTGTTGTAAG
25 540
TGTGAAGCCA GAATTGAGCT AGTAGTAGAA AGCTCAGCAG ACGACCTTCG AGCATTCCAG
600
CAGCTGTTTC TGAACACCCT GTCCTTTGTG TGTCCGTGGT GTGCATCCCA GCAGACTAGT
660
GGCCACCATC ACCATCACCA TTAA
30 684

```

(2) INFORMATION FOR SEQ ID NO:19:

35 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 228 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 40 Protein D 1/3 E7 mutated HPV 18

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

```

45 Met Asp Pro Ser Ser His Ser Ser Asn Met Ala Asn Thr Gln Met Lys
1 5 10 15
Ser Asp Lys Ile Ile Ile Ala His Arg Gly Ala Ser Gly Tyr Leu Pro
20 25 30
Glu His Thr Leu Glu Ser Lys Ala Leu Ala Phe Ala Gln Gln Ala Asp
35 40 45
50 Tyr Leu Glu Gln Asp Leu Ala Met Thr Lys Asp Gly Arg Leu Val Val
55 60
Ile His Asp His Phe Leu Asp Gly Leu Thr Asp Val Ala Lys Lys Phe
65 70 75 80
Pro His Arg His Arg Lys Asp Gly Arg Tyr Tyr Val Ile Asp Phe Thr
85 90 95
Leu Lys Glu Ile Gln Ser Leu Glu Met Thr Glu Asn Phe Glu Thr Met
100 105 110
Ala Met His Gly Pro Lys Ala Thr Leu Gln Asp Ile Val Leu His Leu
115 120 125
60 Glu Pro Gln Asn Glu Ile Pro Val Asp Leu Leu Gly His Gln Gln Leu
130 135 140
Ser Asp Ser Glu Glu Glu Asn Asp Glu Ile Asp Gly Val Asn His Gln
145 150 155 160
65 His Leu Pro Ala Arg Arg Ala Glu Pro Gln Arg His Thr Met Leu Cys
165 170 175

```

Met Cys Cys Lys Cys Glu Ala Arg Ile Gln Leu Val Val Glu Ser Ser  
 180 185 190  
 Ala Asp Asp Leu Arg Ala Phe Gln Gln Leu Phe Leu Asn Thr Leu Ser  
 195 200 205  
 5 Phe Val Cys Pro Trp Cys Ala Ser Gln Gln Thr Ser Gly His His His  
 210 215 220  
 His His His  
 225

10 (2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 837 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 Protein D 1/3 E6 - His HPV 18

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

20 ATGGATCCAA GCAGCCATTC ATCAAATATG GCGAATACCC AAATGAAATC AGACAAAATC  
 60 ATTATTGCTC ACCGTGGTGC TAGCGGTTAT TTACCAGAGC ATACGTTAGA ATCTAAAGCA  
 120 CTTGCGTTTG CACAACAGGC TGATTATTTA GAGCAAGATT TAGCAATGAC TAAGGATGGT  
 25 180 CGTTTAGTGG TTATTCACGA TCACTTTTGA GATGGCTTGA CTGATGTTGC GAAAAAATTC  
 240 CCACATCGTC ATCGTAAAGA TGGCCGTTAC TATGTCATCG ACTTTACCTT AAAAGAAATT  
 30 300 CAAAGTTTAG AAATGACAGA AAACCTTTGAA ACCATGGCGC GCTTTGAGGA TCCAACACGG  
 360 CGACCTACA AGCTACCTGA TCTGTGCACG GAACTGAACA CTTCACTGCA AGACATAGAA  
 420 ATAACCTGTG TATATTGCAA GACAGTATTG GAACTTACAG AGGTATTTGA ATTTGCATTT  
 35 480 AAAGATTTAT TTGTGGTGTA TAGAGACAGT ATACCGCATG CTGCATGCCA TAAATGTATA  
 540 GATTTTATT CTAGAATTAG AGAATTAAGA CATTATTCAG ACTCTGTGTA TGGAGACACA  
 40 600 TTGGAAAAAC TAACTAACAC TGGGTATAC AATTTATTAA TAAGGTGCCT GCGGTGCCAG  
 660 AAACCGTTGA ATCCAGCAGA AAAACTTAGA CACCTTAATG AAAACGACG ATTCACAAC  
 720 ATAGCTGGGC ACTATAGAGG CCAGTGCCAT TCGTGCTGCA ACCGAGCACG ACAGGAACGA  
 45 780 CTCCAACGAC GCAGAGAAAC ACAAGTAACT AGTGGCCACC ATCACCATCA CCATTAA  
 837

50 (2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

55 (A) LENGTH: 279 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 Protein D 1/3 E6 - His HPV 18

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

60 Met Asp Pro Ser Ser His Ser Ser Asn Met Ala Asn Thr Gln Met Lys  
 1 5 10 15  
 Ser Asp Lys Ile Ile Ala His Arg Gly Ala Ser Gly Tyr Leu Pro  
 20 25 30  
 65 Glu His Thr Leu Glu Ser Lys Ala Leu Ala Phe Ala Gln Ala Asp  
 35 40 45

Tyr Leu Glu Gln Asp Leu Ala Met Thr Lys Asp Gly Arg Leu Val Val  
 50 55 60  
 Ile His Asp His Phe Leu Asp Gly Leu Thr Asp Val Ala Lys Lys Phe  
 65 70 75 80  
 5 Pro His Arg His Arg Lys Asp Gly Arg Tyr Tyr Val Ile Asp Phe Thr  
 85 90 95  
 Leu Lys Glu Ile Gln Ser Leu Glu Met Thr Glu Asn Phe Glu Thr Met  
 100 105 110  
 10 Ala Arg Phe Glu Asp Pro Thr Arg Arg Pro Tyr Lys Leu Pro Asp Leu  
 115 120 125  
 Cys Thr Glu Leu Asn Thr Ser Leu Gln Asp Ile Glu Ile Thr Cys Val  
 130 135 140  
 Tyr Cys Lys Thr Val Leu Glu Leu Thr Glu Val Phe Glu Phe Ala Phe  
 145 150 155 160  
 15 Lys Asp Leu Phe Val Val Tyr Arg Asp Ser Ile Pro His Ala Ala Cys  
 165 170 175  
 His Lys Cys Ile Asp Phe Tyr Ser Arg Ile Arg Glu Leu Arg His Tyr  
 180 185 190  
 20 Ser Asp Ser Val Tyr Gly Asp Thr Leu Glu Lys Leu Thr Asn Thr Gly  
 195 200 205  
 Leu Tyr Asn Leu Leu Ile Arg Cys Leu Arg Cys Gln Lys Pro Leu Asn  
 210 215 220  
 Pro Ala Glu Lys Leu Arg His Leu Asn Glu Lys Arg Arg Phe His Asn  
 225 230 235 240  
 25 Ile Ala Gly His Tyr Arg Gly Gln Cys His Ser Cys Cys Asn Arg Ala  
 245 250 255  
 Arg Gln Glu Arg Leu Gln Arg Arg Arg Glu Thr Gln Val Thr Ser Gly  
 260 265 270  
 30 His His His His His His  
 275

## (2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1152 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

Protein D1/3 E6 E7 His/ HPV 18  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

ATGGATCCAA GCAGCCATTC ATCAAATATG GCGAATACCC AAATGAAATC AGACAAAATC  
 60  
 45 ATTATTGCTC ACCGTGGTGC TAGCGGTTAT TTACCAGAGC ATACGTTAGA ATCTAAAGCA  
 120  
 CTTGCGTTTG CACAACAGGC TGATTATTTA GAGCAAGATT TAGCAATGAC TAAGGATGGT  
 180  
 CGTTTAGTGG TTATTCACGA TCACTTTTGA GATGGCTTGA CTGATGTTGC GAAAAAATTC  
 240  
 50 CCACATCGTC ATCGTAAAGA TGGCCGTTAC TATGTCATCG ACTTTACCTT AAAAGAAATT  
 300  
 CAAAGTTTAG AAATGACAGA AAACCTTGAA ACCATGGCGC GCTTTGAGGA TCCAACACGG  
 360  
 55 CGACCCTACA AGCTACCTGA TCTGTGCACG GAACTGAACA CTTCACTGCA AGACATAGAA  
 420  
 ATAACCTGTG TATATTGCAA GACAGTATTG GAACTTACAG AGGTATTTGA ATTTGCATTT  
 480  
 AAAGATTTAT TTGTGGTGTA TAGAGACAGT ATACCGCATG CTGCATGCCA TAAATGTATA  
 540  
 60 GATTTTATT CTAGAATTAG AGAATTAAGA CATTATTCAG ACTCTGTGTA TGGAGACACA  
 600  
 TTGGAACAAAC TAACTAACAC TGGGTATAC AATTATTAA TAAGGTGCCT GCGGTGCCAG  
 660  
 65 AAACCGTTGA ATCCAGCAGA AAAACTTAGA CACCTTAATG AAAACGACG ATTTACAAC  
 720

ATAGCTGGGC ACTATAGAGG CCAGTGCCAT TCGTGCTGCA ACCGAGCAGC ACAGGAACGA  
 780  
 CTCCAACGAC GCAGAGAAAC ACAAGTAATG CATGGACCTA AGGCAACATT GCAAGACATT  
 840  
 5 GTATTGCATT TAGAGCCCCA AAATGAAATT CCGGTTGACC TTCTATGTCA CGAGCAATTA  
 900  
 AGCGACTCAG AGGAAGAAAA CGATGAAATA GATGGAGTTA ATCATCAACA TTTACCAGCC  
 960  
 CGACGAGCCG AACCACAACG TCACACAATG TTGTGTATGT GTTGTAAAGT TGAAGCCAGA  
 10 1020  
 ATTGAGCTAG TAGTAGAAAG CTCAGCAGAC GACCTTCGAG CATTCCAGCA GCTGTTTCTG  
 1080  
 AACACCCTGT CCTTGTGTG TCCGTGGTGT GCATCCCAGC AGACTAGTGG CCACCATCAC  
 1140  
 15 CATCACCATT AA  
 1152

(2) INFORMATION FOR SEQ ID NO:23:

20 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 384 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 25 Protein D1/3 E6 E7 His/ HPV 18

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

30	Met	Asp	Pro	Ser	Ser	His	Ser	Ser	Asn	Met	Ala	Asn	Thr	Gln	Met	Lys
	1				5					10					15	
	Ser	Asp	Lys	Ile	Ile	Ile	Ala	His	Arg	Gly	Ala	Ser	Gly	Tyr	Leu	Pro
				20					25					30		
	Glu	His	Thr	Leu	Glu	Ser	Lys	Ala	Leu	Ala	Phe	Ala	Gln	Ala	Asp	
				35				40					45			
35	Tyr	Leu	Glu	Gln	Asp	Leu	Ala	Met	Thr	Lys	Asp	Gly	Arg	Leu	Val	Val
		50				55						60				
	Ile	His	Asp	His	Phe	Leu	Asp	Gly	Leu	Thr	Asp	Val	Ala	Lys	Lys	Phe
	65					70					75					80
	Pro	His	Arg	His	Arg	Lys	Asp	Gly	Arg	Tyr	Tyr	Val	Ile	Asp	Phe	Thr
					85					90					95	
40	Leu	Lys	Glu	Ile	Gln	Ser	Leu	Glu	Met	Thr	Glu	Asn	Phe	Glu	Thr	Met
				100					105					110		
	Ala	Arg	Phe	Glu	Asp	Pro	Thr	Arg	Arg	Pro	Tyr	Lys	Leu	Pro	Asp	Leu
				115				120					125			
45	Cys	Thr	Glu	Leu	Asn	Thr	Ser	Leu	Gln	Asp	Ile	Glu	Ile	Thr	Cys	Val
		130				135						140				
	Tyr	Cys	Lys	Thr	Val	Leu	Glu	Leu	Thr	Glu	Val	Phe	Glu	Phe	Ala	Phe
	145					150					155					160
	Lys	Asp	Leu	Phe	Val	Val	Tyr	Arg	Asp	Ser	Ile	Pro	His	Ala	Ala	Cys
				165						170					175	
50	His	Lys	Cys	Ile	Asp	Phe	Tyr	Ser	Arg	Ile	Arg	Glu	Leu	Arg	His	Tyr
				180					185					190		
	Ser	Asp	Ser	Val	Tyr	Gly	Asp	Thr	Leu	Glu	Lys	Leu	Thr	Asn	Thr	Gly
				195				200					205			
55	Leu	Tyr	Asn	Leu	Leu	Ile	Arg	Cys	Leu	Arg	Cys	Gln	Lys	Pro	Leu	Asn
		210				215						220				
	Pro	Ala	Glu	Lys	Leu	Arg	His	Leu	Asn	Glu	Lys	Arg	Arg	Phe	His	Asn
	225					230					235					240
	Ile	Ala	Gly	His	Tyr	Arg	Gly	Gln	Cys	His	Ser	Cys	Cys	Asn	Arg	Ala
				245						250					255	
60	Arg	Gln	Glu	Arg	Leu	Gln	Arg	Arg	Arg	Glu	Thr	Gln	Val	Met	His	Gly
				260					265					270		
	Pro	Lys	Ala	Thr	Leu	Gln	Asp	Ile	Val	Leu	His	Leu	Glu	Pro	Gln	Asn
				275				280					285			
65	Glu	Ile	Pro	Val	Asp	Leu	Leu	Cys	His	Glu	Gln	Leu	Ser	Asp	Ser	Glu
		290					295						300			

	Glu	Glu	Asn	Asp	Glu	Ile	Asp	Gly	Val	Asn	His	Gln	His	Leu	Pro	Ala
	305					310					315					320
	Arg	Arg	Ala	Glu	Pro	Gln	Arg	His	Thr	Met	Leu	Cys	Met	Cys	Cys	Lys
					325					330						335
5	Cys	Glu	Ala	Arg	Ile	Glu	Leu	Val	Val	Glu	Ser	Ser	Ala	Asp	Asp	Leu
				340					345					350		
	Arg	Ala	Phe	Gln	Gln	Leu	Phe	Leu	Asn	Thr	Leu	Ser	Phe	Val	Cys	Pro
			355					360					365			
10	Trp	Cys	Ala	Ser	Gln	Gln	Thr	Ser	Gly	His	His	His	His	His	His	
	370						375						380			

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : A61K 39/12, 39/39 // C07K 19/00, C12N 15/62	A3	(11) International Publication Number: <b>WO 99/33868</b>  (43) International Publication Date: 8 July 1999 (08.07.99)
(21) International Application Number: PCT/EP98/08563 (22) International Filing Date: 18 December 1998 (18.12.98) (30) Priority Data: 9727262.9                      24 December 1997 (24.12.97)    GB (71) Applicant (for all designated States except US): SMITHK- LINE BEECHAM BIOLOGICALS S.A. [BE/BE]; Rue de l'Institut 89, B-1330 Rixensart (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): DALEMANS, Wilfried, L., J. [BE/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE). GERARD, Catherine, Marie, Ghislaine [BE/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE). (74) Agent: DALTON, Marcus, Jonathan, William; SmithKline Beecham, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims</i> <i>and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 16 September 1999 (16.09.99)
(54) Title: HUMAN PAPILLOMAVIRUS VACCINE (57) Abstract <p>The present invention provides Human Papilloma Virus (HPV) fusion proteins, linked to an immunological fusion partner that provides T helper epitopes to the HPV antigen. Vaccine formulations are provided that are useful in the treatment or Prophylaxis of HPV induced tumours.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/08563

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K39/12 A61K39/39 //C07K19/00,C12N15/62

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 96 19496 A (CSL LIMITED ET AL.) 27 June 1996 (1996-06-27) cited in the application page 7, line 6 - line 11; claims --- -/--	1-15

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

15 July 1999

Date of mailing of the international search report

22/07/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Ryckebosch, A

# INTERNATIONAL SEARCH REPORT

In. .ational Application No

PCT/EP 98/08563

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>R.S. CHU ET AL.: "CpG OLIGODEOXYNUCLEOTIDES ACT AS ADJUVANTS THAT SWITCH ON T HELPER 1 (Th1) IMMUNITY." JOURNAL OF EXPERIMENTAL MEDICINE, vol. 186, no. 10, 1 November 1997 (1997-11-01), pages 1623-1631, XP002910130 NEW YORK, N.Y., US ISSN: 0022-1007</p> <p>page 1624, right-hand column, line 1; figure 1; table 1</p> <p>page 1629, right-hand column, paragraph 4</p> <p>- page 1630, right-hand column, paragraph 1</p>	1,3-15
Y	<p>WO 97 01640 A (SMITHKLINE BEECHAM BIOLOGICALS S.A.)</p> <p>16 January 1997 (1997-01-16)</p> <p>cited in the application</p> <p>page 2, line 3 - line 10; claims 1-8</p> <p>page 5, line 8 - line 19</p>	2
A	<p>WO 91 18926 A (A. FORSGREN)</p> <p>12 December 1991 (1991-12-12)</p> <p>page 2, line 31 - line 34; claims 1,12,13,16</p> <p>page 5, line 22 - line 35</p>	1,2

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 98/08563

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 13 and 14 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/08563

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9619496 A	27-06-1996	AU 693627 B	02-07-1998
		AU 4322996 A	10-07-1996
		CA 2207741 A	27-06-1996
		EP 0796273 A	24-09-1997
		JP 10510989 T	27-10-1998
		ZA 9510832 A	04-07-1996
WO 9701640 A	16-01-1997	AU 6304996 A	30-01-1997
		CA 2222456 A	16-01-1997
		CZ 9704223 A	17-06-1998
		EP 0835318 A	15-04-1998
		NO 976060 A	17-02-1998
		PL 324906 A	22-06-1998
WO 9118926 A	12-12-1991	SE 466259 B	20-01-1992
		AT 170531 T	15-09-1998
		AU 650011 B	09-06-1994
		AU 7559391 A	31-12-1991
		CA 2083172 A	01-12-1991
		DE 69130116 D	08-10-1998
		DE 69130116 T	18-02-1999
		EP 0594610 A	04-05-1994
		ES 2119776 T	16-10-1998
		FI 925460 A	30-11-1992
		SE 9001949 A	01-12-1991
		US 5888517 A	30-03-1999
		US 5858677 A	12-01-1999